

This Page Is Inserted by IFW Operations
and is not a part of the Official Record

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents *will not* correct images,
please do not report the images to the
Image Problem Mailbox.

(19)



Europäisches Patentamt
European Patent Office
Office européen des brevets

(11) Publication number:

**0 260 404
A2**

(12)

EUROPEAN PATENT APPLICATION

(21) Application number: 87110370.1

(22) Date of filing: 17.07.87

(51) Int. Cl. 4: C12N 15/00, C12P 21/02,
C12N 9/30, C12N 9/34,
C12P 7/06, C12P 1/00,
C12C 11/04, C12N 1/16,
A21D 8/04, C12N 1/18

The microorganism(s) has (have) been deposited with American Type Culture Collection under number(s) DSM 3794; DSM 3792; DSM 3793; DSM 3795; DSM 3796; DSM 3797; DSM 3820; DSM 3798; DSM 3799; DSM 3788; DSM 3789; DSM 3790; DSM 3791.

(30) Priority: 21.08.86 EP 86111586

(31) Date of publication of application:
23.03.88 Bulletin 88/12

(84) Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI LU NL SE

(71) Applicant: Heineken Technisch Beheer B.V.
Burg. Smeetsweg 1
NL-2382 PH Zoeterwoude(NL)

(72) Inventor: Strasser, Alexander
Zonser Strasse 6
D-4000 Düsseldorf(DE)
Inventor: Martens, Feodor Bernard
Hogenwoerd 13
NL-2311 HE Leiden(NL)
Inventor: Dohmen, Jürgen
Bretonenstrasse 20
D-4005 Meerbusch 1(DE)
Inventor: Hollenberg, Cornelius P.
Chopinstrasse 7
D-4000 Düsseldorf(DE)

(74) Representative: Patentanwälte Grünecker,
Kinkeldey, Stockmair & Partner
Maximilianstrasse 58
D-8000 München 22(DE)


(54) Amylolytic enzymes producing microorganisms constructed by recombinant DNA technology and their use for fermentation processes.

(57) This invention provides a method for producing amylolytic enzymes by culturing a microorganism, having received as a result of recombinant DNA technology DNA sequences from a donor yeast comprising the coding sequences for the amylolytic enzymes wherein the host microorganism is capable of expressing said amylolytic enzymes. Furthermore, this invention provides microorganisms genetically engineered as to being able to produce and express the amylolytic enzymes, a vector containing the DNA sequences, coding for the amylolytic enzymes and the respective DNA sequences. The said host microorganisms are useful in the production of biomass and many fermentation processes, preferably in the production of special beers.

EP 0 260 404 A2

0 260 404

FIG. 1



La

AMYLOLYTIC ENZYMES PRODUCING MICROORGANISMS, CONSTRUCTED BY RECOMBINANT DNA TECHNOLOGY AND THEIR USE FOR FERMENTATION PROCESSES

Species of the yeast Schwanniomyces hydrolyse starch to glucose as the result of the production of two extracellular enzymes, namely amylase (α -1,4-glucan 4-glucanohydrolase E.C.3.2.1.1) and glucoamylase (syn. amyloglucosidase) (α -1,4-glucanglucohydrolase, E.C.3.2.1.3. with debranching activity E.C.3.2.1.9.).

5 The amylolytic systems of Schwanniomyces castelli and Schwanniomyces alluvius are well documented (Oteng-Gyang et al., 1981; Sills et al., 1982, 1984a, 1984b; Wilson et al., 1982).

It is desirable to work with Schwanniomyces spp. because the α -amylase has the liquefying activity and the glucoamylase has the debranching activity, and because the amylolytic system is sensitive to pasteurisation conditions normally employed in brewing. (Sills et al., 1984a and b).

10 The yeast Saccharomyces, having none of the mentioned activities of the yeast Schwanniomyces, is the mostly used organism in brewery, baking processes and in the ethanol-fermentation industry. In the fermentation processes, starch is conventionally used as a raw material. Since, however, as mentioned, Saccharomyces cerevisiae is not able to hydrolyse starch, the traditional conversion of starch into ethanol and CO₂ by Saccharomyces cerevisiae requires two enzymatic degradation processes prior to the
15 fermentation activity by Saccharomyces cerevisiae, namely liquefaction, carried out by adding α -amylase and saccharification by adding glucoamylase with debranching activity (Fogarty et al., 1979).

In the European patent application no. 0 125 615, the use of yeast of the genus Schwanniomyces for the production of low calorie beers has already been described. Using yeast of the genus Schwanniomyces provides an organism in the fermentation procedures which is able to produce the amylolytic enzymes, and
20 therefore the mentioned enzymes do not have to be added prior to the fermentation procedure. Nevertheless, the yeast Schwanniomyces is not at all as efficient and adapted as Saccharomyces cerevisiae in brewery and industrial ethanol production. In the above mentioned European patent application, it has therefore been proposed to use Schwanniomyces in addition to the more efficient genus Saccharomyces, the latter still being the essential microorganism for the fermentation of degraded carbohydrates to ethanol
25 and the starch degrading enzymes from Schwanniomyces. The enzymes are still added separately as usual in the state of the art.

Several amylolytic enzymes encoding genes could be cloned and expressed in Saccharomyces cerevisiae, namely the α -amylase gene of mice (Thomsen, 1983) and of wheat (Rothstein et al., 1984) and the glucoamylase gene of Saccharomyces diastaticus (Yamashita and Fukui, 1983), of Aspergillus niger -
30 (Nunnberg et al., 1984; Innis et al., 1985), Candida albicans (Cohen et al. 1985) and Rhizopusoryzae European Patent Application 85 115 910.3).

The amylolytic enzymes produced by the above mentioned donor organisms are, however, not useful for the purposes of the present invention. For the use of genetically manipulated Saccharomyces cerevisiae in brewing and baking procedures and the production of industrial ethanol, it is highly desirable to clone the
35 respective genes derived from very closely related organisms, since the production and application of the expressed enzymes is much more acceptable for the food industry and has the required enzymological and chemical properties for industrial processes. Therefore, the amylolytic enzymes of the microorganism Schwanniomyces, which can be produced by recombinant DNA technology in a number of host microorganisms, are more suitable for a number of applications due to the close relationship of the microorganism
40 Schwanniomyces and the host microorganisms used, and due to the low temperature optimum, the temperature sensitivity, and the high stability of the amylolytic enzymes produced by Schwanniomyces.

The α -amylase gene of Schwanniomyces is the first α -amylase gene cloned from a yeast species and is, as a gene from a narrowly related yeast organism, more compatible with the fermentation processes required in the food industry, for example the brewing and baking processes. In addition, compared to other
45 α -amylases, the α -amylase of Schwanniomyces has an optimal enzyme activity at a lower temperature of about 37°C, which allows application at lower fermentation or process temperatures. In addition, the α -amylase of Schwanniomyces can be inactivated at 50°C.

The glucoamylase of Schwanniomyces has a special debranching activity and is hence able to hydrolyse starch virtually completely into glucose, whereas the glucoamylase of Candida albicans or
50 Saccharomyces diastaticus has no debranching activity, and hence are not able to hydrolyse starch completely into glucose. The glucoamylase of Schwanniomyces, besides its specially suited enzymological

and chemical properties, including optimum of pH and temperature, has the additional advantages over the glucoamylases encoded by other genes already cloned in that this enzyme originates from a yeast species and can be heat inactivated at pasteurisation conditions (Sills et al., 1983) which are normally employed in the brewing process.

5 It would therefore be most advantageous for the brewing process, the production of industrial ethanol and yeast biomass to provide cultures of Saccharomyces cerevisiae with its high CO₂ production, its good fermentation efficiency, its high ethanol tolerance, with the ability to hydrolyse starch entirely by synthesising and secreting α -amylase and glucoamylase. Also for the CO₂ production in baking, amylolytic enzyme producing strains can be of great advantage.

10 It was therefore an object of the present invention to provide a microorganism suitable for brewing and baking processes, the production of industrial ethanol and of biomass having the ability to hydrolyse starch entirely if required by being capable of synthesising and secreting α -amylase and glucoamylase.

The desired microorganisms can be provided by genetical manipulation of the microorganisms used by recombinant DNA technology. Recombinant DNA technology is used to clone the commercially interesting 15 genes coding for α -amylase and glucoamylase, whereby DNA fragments are cloned which contain the α -amylase and/or glucoamylase genes of a yeast of the genus Schwanniomyces. After transformation of a host microorganism, the cloned genes coding for amylolytic enzymes are expressed in a host microorganism, for example, Saccharomyces cerevisiae and the gene products are secreted into the medium. The enzymatic pretreatment of starch and/or non-fermentable dextrans, which was necessary to provide the 20 fermenting microorganisms with fermentable sugar, can therefore be avoided, with all advantages with respect to a less expensive and a simplified fermentation processes, for example, in the fields of brewing, production of industrial ethanol and production of biomass. For the baking process, amylolytic yeast can be of great advantage for the preparation of certain types of doughs.

One of the advantages of providing a fermenting microorganism, for example Saccharomyces 25 cerevisiae, having the ability to produce α -amylase and/or glucoamylase is based on the fact that the starch can be continuously and quantitatively degraded, and the biomass or ethanol is simultaneously produced by aerobic or anaerobic conversion of the low molecular carbohydrates resulting from the degradation of the starch. In the case of the above mentioned brewing processes, the result of this complete degradation of the starch is a low calorie beer, since the resulting low molecular carbohydrates are easily fermented by the 30 genetically manipulated microorganisms into ethanol and CO₂.

A further example for the use of amylolytic Saccharomyces cerevisiae strains in beer brewing is the production of special beers like low carbohydrate beer, diet beer and special flavoured beer. In addition these strains allow a broadening of the range of starch-containing raw materials for beer brewing.

Although the general idea of the present invention described above comprises the use of various host 35 microorganisms which are suitable in one of the afore mentioned fermentation processes or the production of biomass, a preferred example of constructing a microorganism so as to be able to express and secrete amylolytic enzymes is now described.

For the construction of a yeast of the species Saccharomyces cerevisiae, a transformation system with a particular marker was developed, providing both regulation of the amylase genes and over-production of 40 their gene product in the yeast cells.

According to the present invention, in addition a shuttle cosmid vector has been constructed which can be used as a shuttle vector for transformation of Schwanniomyces, Saccharomyces, Escherichia coli and Schizosaccharomyces pombe.

According to a preferred embodiment of the present invention, the yeast Schwanniomyces castellii acts 45 as a donor for the cloning of α -amylase and/or glucoamylase genes. The yeast Saccharomyces cerevisiae acts as a host yeast being capable of expressing the mentioned genes for the amylolytic enzymes after having received the cloned genes adapted for expression by transformation procedures and further being capable of secreting the gene products.

As a result of genetic manipulation, furthermore, an over-production of the amylolytic enzymes in the 50 yeast Schwanniomyces aluvius is possible. This over-production of amylolytic enzymes in Schwanniomyces strains are of great advantage for the production of these enzymes as such, which after recovering in a known manner, can still be used as described above in fermentation processes in which it is still necessary to add the amylolytic enzymes separately.

The new Schwanniomyces vectors control the function of replication and maintenance in Schwanniomyces. 55 This replication sequence (SwARS1) isolated from Schwanniomyces chromosomal DNA is responsible for autonomous replication in, and high frequency transformation of Schwanniomyces. The SwARS1-element is also functional in Saccharomyces cerevisiae and hence leads to high frequency transformation of Saccharomyces cerevisiae mutants with SwARS1-plasmid.

One suitable representative of Schwanniomyces vectors is pCJD5-1, a hybrid plasmid composed of the SwARS1 sequence, the cos (cohesive) sequence of bacteriophage λ , the pBR322 sequence and the TRP5 gene of Saccharomyces cerevisiae used as a selectable marker.

This plasmid is a new Saccharomyces cerevisiae -Schwanniomyces alluvius -Escherichia coli cosmid shuttle vector with suitable cloning sites for the construction of genomic DNA libraries.

The present invention provides furthermore a new simple and quick transformation procedure for the yeasts Schwanniomyces alluvius, Saccharomyces cerevisiae, Schizosaccharomyces pombe, Kluyveromyces lactis.

Using the above vector and transformation system it is possible to provide new genetically engineered yeast strains of the genus Saccharomyces cerevisiae which are capable of synthesising amylases in culture for mass production.

A consequence of this mass production will be the use of this modified Saccharomyces cerevisiae strain in aerobic and anaerobic starch converting processes to yield yeast biomass, ethanol and CO₂.

According to this invention, new DNA sequences are provided which lead to expression and secretion of proteins in Saccharomyces cerevisiae, Schwanniomyces alluvius, Schizosaccharomyces pombe and Kluyveromyces lactis, since the new DNA sequences contain new secretion signals which allow secretion in the yeasts named.

In addition, sequences are provided by the new DNA sequences which allow regulated gene expression.

Another object of this invention is the provision of a vector without any E. coli sequence for the integration of the amylase genes into the Saccharomyces cerevisiae genome.

Detailed Description of the Present Invention

In the following, a more detailed description of all aspects of the present invention is given.

BRIEF DESCRIPTION OF THE FIGURES

Figure 1 :

A: Alignment of plasmid pYc1- α derived by insertion of a 5 kb fragment of S. castellii DNA, which contains the α -amylase gene, into the cosmid pYc1.

B: Restriction map of the cloned EcoRI fragment of S. castellii.

C: Fragments (1, 2 and 3) subcloned into the plasmid pJDB207 do not lead to functional expression of the α -amylase gene in S. cerevisiae GRF18 transformants.

D: Sequencing strategy:

For sequencing according to Maxam & Gilbert (1980) the EcoRI/Sal1 fragments were 3' labelled with γ -³²P-dTTP. The orientation and the extension of the sequencing are indicated by arrows.

Figure 2:

The nucleotide sequence of the 5' end of the α -amylase structural gene and its 5' flanking region.

Figure 3:

Amino acid sequences of 5 isolated tryptic peptides of glucoamylase.

Figure 4:

Secreted α -amylase activity of the transformants in different growth phases.

The cells were grown up overnight in precultures with 0.67 % YNB and 4 % glucose. After washing the cells with the medium of the different main cultures, the main cultures were inoculated to 0.D.₆₀₀ of 0.2. The main cultures (1 liter YNB medium with the given carbon sources in Fernbach vessels) of the *S. alluvius* NGA23 transformants were buffered with 0.2 M sodium phosphate buffer pH 6.2 and of the *S. cerevisiae* MC34 transformants with 0.1 M citrate buffer pH 6.2. The cultures were incubated at 30°C with constant shaking (2 Hertz) on a longitudinal shaker. The measurements of the α -amylase activity and the glucos concentration in the culture supernatant were determined as described. For further details, see specification.

Figure 5:

FPLC ion exchange chromatography on a Mono Q column for separation of the α -amylase protein from concentrated extracellular medium of *S. cerevisiae* MC34 transformed with pYc1- α .

Using a 10 mM Tris-Cl buffer pH 7.5, the α -amylase could be eluted by an increased NaCl gradient at approximately 300 mM NaCl.

a: A_{284} nm

b: % buffer B

Buffer A: 10 mM Tris-Cl, pH 7.5

Buffer B: 10 mM Tris-Cl, pH 7.5, 2 M NaCl

Figure 6:

Analysis of the purified α -amylase protein from extracellular medium of *S. cerevisiae* MC34 transformed with the plasmid pYc1- α by SDS-PAGE

Lane a: purified α -amylase

Lane b: molecular weight standard proteins

myosin from rabbit muscle (205 kd); β -galactosidase from *E. coli* (116 kd); phosphorylase b from rabbit muscle (97 kd); albumin, bovine (66 kd); albumin, egg (45 kd); carbonic anhydrase from bovine erythrocytes (29 kd).

Figure 7 :

Thin layer chromatography of starch hydrolysis products.

The α -amylase activity of culture supernatants on soluble starch after 2 hours at 40°C was additionally characterized and determined by a thin layer chromatography of the reaction products using Kies Igel 60 from Merck. The chromatography was done with 2:2:1 acetone - isopropanol - 0.1 M lactic acid solvent system. The chromatogram was developed using naphtoresorcinol as described by Touchstone & Dobbins (1978).

Figure 8:

Construction of *S. alluvius* plasmids

A a: The plasmid YRp7HIS4 which was isolated from a genomic Sau3A library of *S. castellii* DNA in the plasmid YRp7 (Struhl et al. 1979) by complementation of the his4-519 mutation of *S. cerevisiae* AH22, was digested with BamHI and ClaI and religated. One of the resulting plasmids (YpRHIS4) contains a 1.5 kb BamHI/ClaI fragment from the his4 region of *S. castellii* in YRp7.

b: Deletion of the 1.45 kb ArS1-TRP1-EcoRI fragment from the plasmid YpRHIS4.

c: Insertion of the 5 kb AMY-EcoRI fragment into the plasmid pBRSwARS1.

d: Isolation of the 3.2 kb TRP5-BamHI fragment from the plasmid pYAS1 by preparative gel electrophoresis.
 e: Insertion of the 3.2 kb TRP5-BamHI fragment into the plasmids YpRHIS4, pBRSwARS1 and pBRSwARS1- α .

B The plasmids YRp7TRP5-A and -B were constructed by inserting the 3.2 kb TRP5-BamHI fragment from pYAS1 into YRp7. Plasmids with both orientations (-A and -B) of the fragment were obtained.

C The plasmids pEK2TRP5-A and -B were constructed by inserting the 3.2 kb TRP5-BamHI fragment from pYAS1 into the *K. lactis* vector pEK2 in both possible orientations (-A and -B).

10 Figure 9:

Copy number estimation of the plasmid YRpJD2 in *S. alluvius* NGA23 and in *S. cerevisiae* X3656 2D.

A Ethidium bromide stained plasmid and yeast minilysate DNA separated in a 0.7 % agarose gel.

15 B Autoradiogram after Southern blot and hybridisation with nick translated YRpJD1 DNA (48 hours exposition at -70°C).

C Autoradiogram of lanes j-q after 312 hours exposition at -70°C

D Map of the hybridisation probe YRpJD1 (thin line: pBR322; open boxes, *S. cerevisiae* DNA; black box, *S. castellii* DNA).

20 Lanes a-c, control plasmids; a: YRpJD1 EcoRI

b: YRpJD1 uncleaved

c: YRpJD2 uncleaved.

Lanes d-g, yeast minilysate DNA; d-f, DNA of *S. alluvius* NGA23: YRpJD2 transformants, d and f, cleaved with EcoRI; e and g, uncleaved; h and i, NGA23:YRpJD1 DNA cleaved with EcoRI (h) and uncleaved (i); j and k, DNA of untransformed NGA23 cleaved with EcoRI (j) and uncleaved (k); l and m, DNA of *S. cerevisiae* X3656 2D : YRpJD2 transformant cleaved with EcoRI (l) and uncleaved (m); n-q, DNA of untransformed X3656 2D cleaved with EcoRI (n and p) and uncleaved (o and q).

30 Figure 10:

Copy number estimation of different plasmids in *S. alluvius* NGA23.

35 A Ethidium bromide stained EcoRI fragments of control plasmids and yeast minilysate DNA after separation in a 0.7 % agarose gel.

B Autoradiogram after Southern blot and hybridisation with nick translated YRp7HIS4 DNA (15 hours exposition at -70°C).

40 C Autoradiogram after 6 hours (lanes a-d) and after 125 hours (lanes e-p) exposition at -70°C respectively (faint bands in lanes j and m and the position of the bands of the chromosomal HIS4 locus are indicated by arrows).

D Scheme of the homologies of YRpJD1 and the chromosomal *Schwanniomyces* HIS4 region to the hybridisation probe YRp7HIS4.

45 lanes a-e, control plasmids: a, pEK2TRP5-A; b, YRp7TRP5-A, c, pYAS1; d, YRpJD1; lane e, DNA of untransformed *S. alluvius* NGA23; lane f, DNA of untransformed *S. castellii* 26076; lanes g-p, DNA of *S. alluvius* NGA23 transformed with : g, YRpJD1; h and j, YRp7TRP5-A; i, k and l, YRp7TRP5-B; m, pEK2TRP5-A; n, pEK2TRP5-B; o and p, pYAS1 (DNA in lane o is uncleaved).

Figure 11:

50

Comparison of the copy number of SwARS1 and ARS1 plasmids in *S. alluvius* and *S. cerevisiae*.

A Ethidium bromide stained EcoRI fragments of control plasmids and yeast minilysate DNA after separation in a 0.7 % agarose gel.

55 B Autoradiogram after Southern blot and hybridisation with nick translated YRpJD2- α DNA. The position of the 2.0 kb band of the chromosomal *S. cerevisiae* TRP5 locus is indicated by an arrow.

lanes a-d, control plasmids: a, pCJD5-1; b, YRpJD2- α ; c, YRpJD1; d, YRp7TRP5-A;
lanes e-h, DNA of *S. alluvius* NJD1; , f and i, untransformed; g, transformed with pCJD5-1; h, transformed
with YRpJD2;
lanes j-m, DNA of *S. alluvius* NGA23; j, untransformed; transformed with: k, YRpJD2- α ; l, YRpJD2; m,
5 YRpJD1;
lanes n-r, DNA of *S. cerevisiae* X3656 2D, transformed with: n, YRpJD2- α ; o, YRpJD2; p, YRpJD1; q,
YRp7TRP5-A; lane r, untransformed.

10 **Figure 12:**

In vivo labelling of extracellular proteins from *S. alluvius* NGA23 transformants.

The transformants were grown under repressed and induced conditions (see Figures 4a, b and c) in 1
15 ml culture after addition of 100 μ Ci 35 S-methionine (800 Ci/mmol). After incubation at 30°C for different
times (18, 22, 28, 43 and 90 hours) 30 μ l of the cell-free supernatant were analysed by SDS-PAGE and
autoradiography.

- a, NGA23:YRpJD2 in YNB, 2 % glucose (repressed);
- b, NGA23:YRpJD2 in YNB, 2 % maltose (induced);
- 20 c, NGA23: YRpJD2- α in YNB, 2 % glucose repressed);
- d, NGA23:YRpJD2- α in YNB, 2 % maltose (induced). The positions of the molecular weight standard
proteins (see Figure 6) are indicated by arrows.

25 **Figure 13:**

Restriction map and restriction analysis of the cosmids pCJD5-1 and pCJD5-2.

- A Restriction map
- 30 B Restriction fragments separated in a 0.7 % agarose gel
- lane a, lambda DNA cleaved with EcoRI and HindIII;
- lane q, lambda DNA cleaved with HindIII;
- from left to right:
- plasmids YRpJD2, pCJD5-1, pCJD5-2:
- 35 lanes b, c, d, cleaved with EcoRI;
- lanes h, i, j, cleaved with BamHI;
- lanes k, l, m, cleaved with BglII;
- lanes n, o, p, cleaved with PvuII.

40 **Figure 14:**

Restriction map of the plasmid pKARS2- α .

- 45 The 5 kb AMY-EcoRI fragment from plasmid pYc1- α was inserted into the single EcoRI site of the
plasmid pKARS2-d.

Figure 15:

- 50 A Restriction map of the two subcloned fragments containing the functional glucoamylase gene
(AMG).
- B The AMG fragments were subcloned into the *S. alluvius*-*S. cerevisiae* vector pCJD5-1 resulting in
pCJD5-AMG1 and pCJD5-AMG2, into the *S. cerevisiae*-*S. pombe* vector pJDB207 resulting in pJDB207-
55 AMG1 and pJDB207-AMG2, and into the *K. lactis* vector pEK2, resulting in PEK2-AMG2.

Figure 16:

Inactivation of glucoamylase at different temperatures.

5

Figure 17

The nucleotide sequence of glucoamylase structural gene and flanking regions.

10

Table 1

15

Comparison of preferred codon usage in *S. cerevisiae* and
Schwanniomyces castellii.

20

	amino acid	preferred codons in <u><i>S. cerevisiae</i></u>	preferred codons in <u><i>S. castellii</i></u>
25	ala	GCU, GCC	GCU, GCG
	ser	UCU, UCC	UCA, UCC, AGC, UCU, AGU (no clear preference)
	thr	ACU, ACC	<u>ACU</u> , ACC, ACA
30	val	GUU, GUC	<u>GUU</u> ; GUA, GUC, GUG
	ile	AUU, AUC	AUU; AUC, AUA, AAU (no clear preference)
	asp	GAC	<u>GAU</u> , GAC
35	phe	UUC	UUC, UUU (no clear preference)
	tyr	UAC	<u>UAU</u> , UAC
	cys	UGU	UGU
40	asn	AAC	<u>AAU</u> , AAC
	his	CAC	<u>CAU</u> , CAC
	glu	GAA	<u>GAA</u> , GAG
	gly	GGU	GGU, GGA
45	gln	CAA	CAA
	lys	AAG	AAA
	pro	CCA	CCA, CCU
50	leu	UUG	UUG, UUA, CUU
	arg	AGA	AGA

55

Table II

Transformation frequencies obtained with the modified method for strains of different yeast genera.

Species	Strain	Plasmid	Replicator	Marker	Transformants per μg DNA
<i>S. alluvius</i>	NGA19	YRpJD1	SwARS1	TRP5	300 - 1000
<i>S. alluvius</i>	NGA23	YRpJD1	SwARS1	TRP5	300 - 1000
<i>S. alluvius</i>	HTR36	YRpJD1	SwARS1	TRP5	300 - 1000
<i>S. cerevisiae</i>	AH22	YRp7HIS4	ARS1	HIS4	500 - 2000
<i>S. cerevisiae</i>	X3656	YRpJD1	ARS1	TRP5	500 - 2000
<i>S. cerevisiae</i>	X3656	pYAS1	2 μm	TRP5	1000 - 10000
<i>S. cerevisiae</i>	GRF18	pYc1	2 μm	HIS3	200 - 1000
<i>K. lactis</i>	SD11	pEK2	KARS2	TRP1	500 - 1000
<i>H. polymorpha</i>	LR9	pHARS1	HARS1	URA3	500 - 1500
<i>S. pombe</i>	leu1-32	pJDB207	2 μm	LEU2	1 - 10

Table III

Transformation frequencies, mitotic stabilities and copy numbers in *S. alluvius* NGA23

Plasmid	Origin of replication	Trp ⁺ transformants per µg plasmid DNA	Trp ⁺ -colony forming cells in selective medium	Trp ⁺ -colony forming cells after 15 generations in selective medium	Average copy number per cell	Copy number per Trp ⁺ -colony forming cell
pYe(trp5)1-53	-	0.1-0.2	6 %	0 %	0.1-0.2	2-3
YRp7-TRP5-A	ARS1	5-20	10 %	1 %	0.2-1	2-10
YRp7-TRP5-B	ARS1	5-20	10-20 %	1 %	0.2-1	1-10
PEK2-TRP5-A	ARS1, KARS2	5-20	9 %	1 %	0.2-0.5	2-5
PEK2-TRP5-B	ARS1, KARS2	5-20	11 %	1 %	0.2-0.5	2-5
pYAS1	2µm	50-100	75-90 %	5-10 %	2-3	2-4
YRpJD1	ARS1, SwARS1	300-1000	85-95 %	10-20 %	5-10	6-12

Table IV

Transformation frequencies, mitotic stabilities and average copy number of the plasmids YRp7-TRp5, YRpJD1 and YRpJD2 in S. alluvius NGA23 and S. cerevisiae X3656.

Plasmid	Origin	Transformants per µg DNA		loss of plasmids after 15 generations in non-selective medium		average copy number per cell	
		S.all. NGA23.	S.cer. X3656	S.all. NGA23	S.cer. X3656	S.all. NGA23	S.cer. X3656
YRp7-TRP5-A	ARS1	5-10	500-2000	99 %	80-90 %	0.2-1	5-10
YRpJD1	ARS1, SwARS1	300-1000	500-2000	80-90 %	80-90 %	5-10	5-10
YRpJD2	SwARS1	300-1000	500-2000	80-90 %	80-90 %	5-10	5-10

The yeast strains used are as follows:

- Schwanniomyces castellii ATCC 26076. (DSM 3794)
 30 Schwanniomyces alluvius NGA23 and NGS1. (DSM 3792 and 3793)
 Kluyveromyces lactis SD11. (DSM 3795)
 Schizosaccharomyces pombe leu1-32, his5-303. (DSM 3796)
 Saccharomyces cerevisiae GRF18 leu2-3, leu2-112, his3-11, his3-15. (DSM 3797)
 Saccharomyces cerevisiae AH22a leu2-3, leu2-112, his4-519, can1. (DSM 3820)
 35 Saccharomyces cerevisiae X3656 2D a, ade, arg4-1, his6, leu1, trp5. (DSM 3798)
 Saccharomyces cerevisiae his3 Mal1⁺ (DSM 3799)

The Escherichia coli strains used are as follows

- Escherichia coli, HB101 F⁻, hsd S20, r_B⁻ m_B⁻, recA13, ara-14, proA2, lacY1, galK2, rpsL20 (Sm^r), xy15, mtl-1, supE44, λ⁻ (DSM3788)
 40 Escherichia coli JA221, recA-1, leuB6, trpE5, hsdR⁻, hsdM⁺, lacY1. (DSM 3789)
 Escherichia coli BHB2688, N205 recA⁻[imm⁴³⁴, cts, b2, red³, Eam4, Sam7]/λ (DSM 3790)
 Escherichia coli BHB2690, N205 recA⁻[imm⁴³⁴, cts, b2, red³m, Dam15, Sam7]/λ (DSM 3791)

The plasmids and cosmids used are as follows:

- Cosmid pYC1.
 45 Plasmid pEK2.
 Plasmid YRP7.
 Plasmid pJDB207.

As far as they are essential for carrying out the present invention, microorganisms as mentioned above have been deposited with the Deutsche Sammlung für Mikroorganismen (DSM).

- 50 A complete medium for yeast strains was YEPD (2% bactopectone, 1% yeast extract, 2% glucose).
 Minimal medium contained 2% glucose and 0.67% yeast nitrogen base without amino acids (Difco) supplemented with the appropriate amino acids.

Schwanniomyces cells were grown in 2% soluble starch instead of glucose.

- 55 Starch fermenting yeast strains were screened for α-amylase and hence halo formation on YNB minimal medium plates containing 1% glucose, 1% soluble starch and 2% agar. The halos were developed by incubating the plates in an iodine saturated chamber for a few seconds.

Bacterial cells were grown in LB medium (0.5% yeast extract, 1% tryptone, 1% NaCl). When required, ampicillin was added to a final concentration of 150 µg per ml, tetracycline to 20 µg per ml, respectively.

Bacterial transformation was carried out as described by R.W. Davis et al. (1980).

Yeast transformations were performed according to the procedure described in this paper.

Throughout this application, various publications are referenced by the name of the author and date of publication within parentheses. Full citations for these references may be found at the end of the specification as an annex, listed in alphabetical order immediately preceding the claims. The disclosures of these publications in their entireties are hereby incorporated by reference into this application in order to more fully describe the state of the art as known to those skilled therein as of the date of the invention described and claimed herein.

Plasmid DNA was purified either by CsCl gradients centrifugation (Maniatis et al., 1982) or by rapid alkaline extraction of plasmid DNA as described by Birnboim and Doly (1979).

Yeast crude extracts were prepared by the method of Schatz (1979).

Yeast miniprepates were prepared by the method of Sherman et al. (1983).

DNA fragments were isolated by "low melting agarose" procedure as described by Gafner et al. (1983).

15

Methods used for carrying out the invention

DNA sequencing was carried out as described by Maxam and Gilbert (1980). DNA was nick translated according to Rigby et al. (1977).

Southern hybridisation was carried out as described by Southern (1975).

Restriction endonucleases, T4 DNA ligase, *E. coli* polymerase I, Klenow polymerase can be purchased from Boehringer (Mannheim) and used as specified by the suppliers. α -³²P dNTPs and (³⁵S) methionine can be purchased for example from Amersham. Zymolyase 5,000 and zymolyase 60,000 can be obtained from Kirin Brew (Japan). Nitrocellulose filters can be obtained from Schleicher and Schüll.

The cosmid library of *Schwanniomyces castellii* DNA was constructed in the *Saccharomyces cerevisiae* shuttle vector pYc1 (Hohn and Hinnen, 1980) and in the *Saccharomyces cerevisiae*-*Schwanniomyces alluvius*-*E. coli* shuttle vector pCJD5-1 constructed according to this invention as described by Ish Horowitz and Burke (1981).

Protein concentration was determined by the method of Bradford (1976) with bovine serum albumin used as a standard. The protein was separated by SDS polyacrylamide gel electrophoresis according to the procedure of Laemmli (1970).

Amylase activity was measured kinetically with an enzymatic colour test, for example, α -amylase Mercko test, from Merck. The rate of formation of 2-chloro-4-nitrophenol is determined by photometry at 405 nm at 37°C in 0.1 M potassium phosphate activity in the sample material.

Enzyme units (U) are defined as the amount of enzyme catalysing the formation of 1 μ mol 2-chloro-4-nitrophenol \times min⁻¹ at 37°C.

Glucoamylase activity was measured by a stop assay method: After incubation of the sample in 10% soluble starch in 0.05 M KH₂PO₄-NaOH (pH 5.0) at 50°C. The amount of glucose produced is determined by the glucose dehydrogenase method (system glucose Merck). The quantity of NADH formed is proportional to the glucose concentration.

The enzyme unit (U) is defined as the amount of enzyme catalysing the formation of 1 μ mol glucose min⁻¹ at 50°C.

Isolated α -amylase from culture supernatant of *Schwanniomyces castellii* forms a halo on agar plates containing soluble starch after staining with iodine. From these observations, the scheme to clone the α -amylase was based on the assumption that *Saccharomyces cerevisiae* cells which form no halo can be converted to halo-forming ones when a plasmid vector containing the gene for a starch hydrolysing enzyme, namely α -amylase, is present. To clone the α -amylase gene, a recombinant cosmid library was constructed from genomic DNA of *Schwanniomyces castellii* ATCC 26076. Using the cosmid cloning technique described above, a genomic library of about 17,000 independent clones could be generated in the vector pYc1 (Hohn and Hinnen, 1980). The cosmid library was screened for inserts containing the α -amylase gene by transforming *Saccharomyces cerevisiae* GRF18 to histidine prototrophy and checking for the ability to degrade starch.

Approximately 4500 transformants of GRF18 were screened for their ability to form halos after iodine staining on plates containing leucine, glucose and soluble starch. 13 halo-forming transformants were obtained. From these transformants, total yeast DNA was isolated and used to transform *E. coli* JA221 cells with selection for ampicillin resistance. Two different recombinant cosmids containing a 32 kb and a 39 kb

fragment of *Schwanniomyces castellii* DNA respectively were obtained. These recombinant cosmids were reintroduced into GRF18. The transformants showed the His⁺ and halo-forming phenotype. In a mitotic stability test (Beggs, 1978) both cosmid markers, the HIS3 and the halo-forming function, were segregated out simultaneously, indicating that both genes were present on the cosmid.

The smaller cosmid containing the 32 kb insert was used to identify the DNA region responsible for the halo-forming phenotype. To achieve this, the cosmid was digested with EcoRI, religated and transformed into JA221. The DNA from ampicillin resistant *E. coli* cells was used to transform GRF18 with selection for histidine prototrophy.

From five halo-forming transformants, total yeast DNA was isolated and used to transform JA221. Plasmid DNA was isolated from ampicillin resistant *E. coli* transformants. Restriction enzyme analysis revealed that each of these plasmids carries a 5 kb EcoRI *Schwanniomyces castellii* DNA fragment in the original vector pYc1. This plasmid is called pYc1- α (Fig. 1A). The restriction map of the 5 kb EcoRI fragment is presented in Figure 1B.

Fragments 1, 2 and 3 (Fig. 1C), subcloned into the vector pJDB207 (Beggs, 1981) and transformed into *Saccharomyces cerevisiae* GRF18 did not result in α -amylase secreting transformants and hence do not contain the functional gene. Therefore, it can be assumed that the functional α -amylase gene spans the HindIII, BglII sites.

Taking into account that the molecular weight of α -amylase is approximately 61,000 daltons (Wilson et al., 1982), it can be predicted that a DNA fragment of approximately 2 kb should contain the functional gene.

To verify this assumption, DNA sequence analysis was performed according to Maxam and Gilbert (1980). For sequencing both EcoRI-Sall fragments of the 5 kb EcoRI fragments were radioactively labelled at the Sall site (Fig. 1D). DNA sequence analysis revealed a 5' noncoding region followed by an open reading frame spanning the Sall site. The DNA sequence is presented in Fig. 2.

The TATA box (Gannon et al., 1979) usually located 25-32 bp upstream and the CAAT box (Benoist et al., 1980) which has been found about 80 bp upstream from the transcription initiation point, are thought to be important for transcription initiation. While the TATA box is present in almost all examined genes, the CAAT box is not always present.

In the regulation region of the α -amylase gene there is a perfect TATA box at -89 and a CAAT box at -115.

In the 5' region of the α -amylase gene, there is a CAAG sequence at position -7. The codon usage of the α -amylase is given in Table I.

35 Expression of the *Schwanniomyces* α -amylase gene and secretion of the protein by *Saccharomyces cerevisiae*

GRF18:pYc1- α transformants are not able to grow on soluble starch as a sole carbon source. This could be shown to be due to a defect in maltose utilisation. Therefore, the plasmid pYc1- α was used to transform *Saccharomyces cerevisiae* MC34 (his3), having a Mal⁺ phenotype with selection for starch degradation. All histidine phototroph transformants were able to grow on starch as a sole carbon source and could form halos surrounding the colony after staining with iodine. For α -amylase activity measurements, MC34:pYc1- α transformants were cultivated with selection for histidine prototrophy in YNB medium containing glucose.

During the growth of *Saccharomyces cerevisiae* in unbuffered YNB medium, the pH drops to approximately 3.5.

The α -amylase becomes inactivated below a pH of 4 and has a pH optimum at about 6.2 (Sills and Stewart, 1984). Therefore, the medium was buffered with 0.1 M citrate buffer pH 6.2.

During growth of MC34:pYc1- α , crude extract and extracellular medium was assayed for α -amylase activity.

α -amylase activity could only be detected in the extracellular culture medium and not in crude extracts, indicating that the α -amylase is secreted. The total activity in the extracellular medium decreases at late logarithmic growth (Fig. 4D). This loss of enzymatic activity could be explained by proteolytic degradation of α -amylase, because proteolytic activities accumulate during the transition from the logarithmic to stationary growth phase (Ferguson et al., 1973). However the stability of α -amylase could not be improved by adding the protease inhibitor PMSF. It can therefore only be concluded that the inactivation is not due to a serin protease.

The total α -amylase activity in the extracellular medium of stationary grown MC34:pYc1- α cells in complete medium YEPD was significantly higher, compared to cells grown in YNB medium. The most likely explanation is the two-fold higher proteolytic activities when Saccharomyces cerevisiae is grown in YNB, compared to growth in YEPD (Tsai et al., 1973).

5 Sometimes enzymes can be stabilized by the addition of substrate (Katsunuma et al., 1972). MC34:pYc1- α transformants which are grown in YNB containing soluble starch as a sole carbon source showed no difference in the α -amylase activity profile during different growth phases, compared to cells grown in glucose as a sole carbon source, indicating that the α -amylase could not be stabilised in the presence of substrate.

10 However, it was observed that MC34:pYc1- α had an initial lag period in its ability to ferment starch either by an induction period at maltose uptake or probably due to a time dependent accumulation of secreted α -amylase into the medium.

From the recent investigations, it could be shown that the α -amylase gene is also expressed and the gene product secreted by Schizosaccharomyces pombe and Kluyveromyces lactis. For this purpose, 15 Schizosaccharomyces pombe was transformed with plasmid pYc1 α (Fig.1) and Kluyveromyces lactis with plasmid pKARS2- α (Fig. 14a). The former transformants could form halos on starch plates after iodine staining. Quantification of the α -amylase system by these two transformed yeast strains is the subject of the present investigation.

Moreover, the α -amylase gene was cloned into a wild type bottom-fermenting brewers' yeast strain of 20 Saccharomyces cerevisiae var. carlsbergensis by transformation with the plasmid containing α -amylase gene (pYc1- α) and screening for α -amylase activity by halo formation on plates containing soluble starch.

25 Isolation and characterisation of the enzymes α -amylase and glucoamylase from Schwanniomyces castellii

Both enzymes were isolated from the medium of a continuous culture of Schwanniomyces castellii.

Medium: YNB + 1% starch; 20 mM PO₄; pH 6.3.

Conditions: temp. = 30°C; PO₂ = 70 %; D = 0.15 h⁻¹.

Yeast cells were separated from the medium by ultrafiltration (UF) and the filtrate was concentrated with 30 an UF-membrane, cut-off 10,000 D. The concentrated filtrate was loaded on a DEAE-Sephacel column in 20 mM phosphate buffer, pH 5.6.

Glucoamylase was eluted from the column between 150-180 mM NaCl, α -amylase was eluted between 180-250 mM NaCl. Fractions containing enzyme activity were collected, dialysed against 100 mM Ammonium acetate buffer pH 6.5 and concentrated by Amicon YM 10 filtration.

35 Aliquots of ± 4 mg of each protein were further purified using HPLC gel permeation chromatography on a TSK 2000 SW column using 100 mM ammonium acetate buffer pH 6.5.

The enzymes purified in this way were tested for homogeneity by SDS-PAGE, resulting in on single protein band for each enzyme.

40 Preparation of glucoamylase peptide fractions

Glucoamylase (50 mg) was incubated with trypsin (1 mg) for 24 hr., 37°C in 50 mM ammonium acetate buffer pH 6.5, resulting in $\pm 30\%$ breakdown of the enzyme. Peptides were separated by means of 45 HPLC on a TSK 2000 SW column and further purified on RPHPLC using C18 and C8 columns. Elution was carried out with a gradient system 0-100% acetonitrile. From 5 peptides, the amino acid sequence was determined (see Fig. 3).

50 Characterisation of glucoamylase

MW: 120,000 D

pH optimum 5.0

temp. optimum 50°C

55 Km: 1.0 - 1.5 % dextrin (Serva)

pl 3.5
glycosilated
activity repressed by glucos
temp. inactivation - see Fig. 16.

5

Characterisation of α -amylase

MW 53,000 D
10 pH optimum 6.0
temp. optimum 37°C
Km: 0.004 - 0.07 % soluble starch

15 Isolation of secreted α -amylase from *Saccharomyces cerevisiae*

The maximal measurable secreted α -amylase activity from MC34:pYc1-transformants growing in YNB medium could be obtained from cells of the early logarithmic phase (Fig. 4D). It is advantageous to isolate the α -amylase from extracellular YNB medium instead of complex medium containing yeast extract and bactopectone, because the initial specific activity (α -amylase activity/protein ratio) is significantly higher in YNB than in YEPD.

In addition, since the amount of secreted α -amylase is directly correlated to the amount of actively growing yeast cells, a stepwise growth of MC34:pYc1- α in YNB (selection on the plasmid), YEPD (optimal growth condition), and finally YNB (secretion of α -amylase from a high amount of actively growing cells) was used for the isolation of secreted α -amylase. In detail, the transformants were grown in YNB medium containing glucose. When an optical density of approximately 4 at 600 nm was reached (late logarithmic growth phase), the cells were harvested by centrifugation and resuspended in prewarmed (30°C) YEPD to a final OD₆₀₀ of 7.5.

After 2 hours growth at 30°C, the cells were sedimented and resuspended in the same volume of prewarmed (30°C) YNB containing glucose and 0.1M citrate buffer pH 6.2. After 4 hours incubation at 30°C (optical density at 600 nm reached a value of 17), the α -amylase was purified from concentrated extracellular medium by means of FPLC ion exchange chromatography using a MonoQ column (Pharmacia). The α -amylase protein could be eluted in 10 mM Tris-Cl, pH 7.5 at 300 mM NaCl using an increasing NaCl gradient (Fig. 5).

35 The active protein fraction was concentrated by lyophilisation and analyzed by SDS polyacrylamide gel electrophoresis (Fig. 6). The presence of only one band with a molecular weight of about 55,000 D showed that the α -amylase protein was highly purified by this procedure.

For enzyme characterisation, the products of starch degradation by the isolated α -amylase were analysed by thin layer chromatography (Fig. 7). The products of starch hydrolysis are maltose and large amounts of maltotriose and higher oligosaccharides. Glucose is not produced in detectable amounts. This is in agreement with the above mentioned observation, that GRF18(Mal⁻):pYc1- α transformants are not able to grow on starch as a sole carbon source.

45 Transformation of *Schwanniomyces alluvius*

One of the prerequisites for the establishment of a transformation system is the availability of an appropriate mutant which can be functionally complemented by a cloned gene.

Various tryptophan auxotrophic *Schwanniomyces alluvius* mutants were analysed for their defect in the L-tryptophan biosynthetic pathway. The mutants were tested for having a defect in the last step of L-tryptophan biosynthesis, namely the condensation reaction of indole and L-serin, which in *Saccharomyces cerevisiae* is catalysed by tryptophan synthetase.

55

A defect in this last step leads to an accumulation of indol (Manney et al., 1969). The tryptophan auxotrophic mutant NGA23 accumulates indole when grown in YNB medium. In addition, compared to other tryptophan auxotrophs, NGA23 cannot grow on YNB containing indole. It was therefore concluded that NGA23 has a mutation in the tryptophan synthetase gene. NGA23 exhibits a low reversion rate (10^{-9}) and is thus suitable for complementation studies with the cloned Saccharomyces cerevisiae TRP5 gene (Carbon et al., 1977) coding for tryptophan synthetase (Walz et al., 1978; Zalkin et al., 1982). The initial intention was to transform NGA23 via integration.

The transformation experiment was based on the assumption that there is no extensive homology between the TRP5 gene of Saccharomyces cerevisiae and the mutated tryptophan synthetase gene of Schwanniomyces alluvius as revealed by Southern hybridisation of nick translated pYAS1 (Strasser, 1982) DNA to chromosomal DNA of Schwanniomyces alluvius.

From a partial Sau3A bank of Schwanniomyces DNA in YRP7, a 9 kb DNA fragment was cloned, resulting in plasmid YRP7HIS4 complementing the his4 mutation of Saccharomyces cerevisiae strain AH22 (Fig. 8).

It was assumed that a 1.5 kb BamHI-ClaI fragment from this HIS4 region of Schwanniomyces castellii should have enough homology to direct integration into the genome of Schwanniomyces alluvius NGA23 via homologous recombination. Therefore, the plasmid YRpJD2 was constructed (Fig. 8), containing the Saccharomyces cerevisiae Trp5 gene as a selectable marker and the 1.5 kb BamHI-ClaI fragment.

YRpJD2 was used to transform NGA23 by the protoplast method described by Beggs (1978) screening for tryptophan prototrophy on YNB plates. Five transformants per μ g plasmid DNA were obtained.

It seemed unlikely that this surprisingly high transformation frequency was due to integration events, taking into account that the plasmid was not linearised prior to transformation, which would have increased the probability for integration (Orr-Weaver et al., 1981) and that the regeneration frequency of Schwanniomyces using the protoplast method was lower than 0.1%.

The transformation of NGA23 with YRpJD2 is based on autonomous replication as was indicated by the instability of the Trp⁺ phenotype after growth of the transformants under non-selective conditions in YEPD. After 15 generations, 80-90 % of the cells had lost the ability to grow on selective medium.

The autonomous replication was ascertained by transforming E. coli cells with yeast miniprepates and retransformation of Schwanniomyces NGA23. In addition, Southern analysis showed the presence of original YRpJD2 sequences (Fig. 9) and thus exclude integration of the TRP5 gene as a cause of the tryptophan prototrophy.

No transformants except for some abortive transformants could be obtained after transformation of NGA23 with pYe(trp5)1-53 (Walz et al, 1978), which does not contain the 1.5 kb fragment of Schwanniomyces castellii indicating that this Schwanniomyces DNA fragment contains a sequence which allows autonomous replication of plasmid YRpJD2 in Schwanniomyces alluvius NGA23.

Therefore, it is concluded that the 1.5 kb DNA fragment of Schwanniomyces castellii in the vector YRpJD2 contains an autonomous replication sequence, namely SwARS1.

40 Transformation procedure

The regeneration of Schwanniomyces cells using the protoplast method (Beggs, 1978) for transformation is approximately 0.1%. Because the regeneration rate or protoplast formation correlates directly with the transformation frequency, attempts were made to change the conditions for protoplast formation using different concentrations of helicase or zymolyase and different incubation times for the transformation of YRpJD2 into NGA23.

Using helicase, no transformants could be obtained. Only by using zymolyase (1 mg/ml) for 30 min at 37°C, with a regeneration rate of 0.1 %, 5 transformants per μ g YRpJD2 plasmid-DNA could be obtained. Shorter incubation resulted in an increase of the regeneration frequency to approximately 0.5%, but no transformants were obtained.

These findings demonstrate that the protoplast formation (and not the regeneration frequency and hence the enzymatic degradation of the cell wall) is the critical step in the transformation of Schwanniomyces using the protoplast method. Therefore, the transformation procedures described by Ito et al. (1983) and Klebe et al. (1983) were tested.

Both procedures do not require the enzymatic degradation of the cell wall. With the lithium sulphate method of Ito et al. (1983), no transformants were obtained. However, using the freeze-thaw method of Klebe et al. (1983) NGA23 could successfully be transformed with YRpJD2 plasmid, yielding reproducibly 50-100 transformants per μ g YRpJD2 plasmid DNA.

Using the method of Klebe et al (1983), log phase yeast (10 ml) was washed with 5 ml of SBEG (1 M sorbitol, 10 mM bicine, pH 8.35, 3% ethylene glycol) by centrifugation at 1000 x g for 3 min at 22°C. All subsequent incubations were carried out at 30°C. The pellet was resuspended in 0.2 ml SBEG; and after 5 min, 1 to 20 µl plasmid DNA was added. After 10 min, the preparation was placed in a -70°C freezer for 10 min (or longer) and thawed by rapid agitation in a 37°C water bath. 1.5 ml of 40% purified PEG 1000 (Klebe et al., 1983), 200 mM bicine pH 8.35 (stored frozen) was added; and after a 1 h incubation, the cells were washed with 2 ml 0.15 mM NaCl, 10 mM bicine pH 8.35. After centrifugation, the pellet was resuspended in 1 ml 0.15 mM NaCl, 10 mM bicine pH 8.35, and plated on selective medium. Using the conditions described above, the plasmid DNA is added before the cells are frozen at -70°C. However, the possibility of adding the DNA to the frozen cells just before thawing would be of advantage to have competent cells at any time for the transformation with any plasmid available.

The attempts of adding YRpJD2 DNA to frozen cells of the *Schwanniomyces alluvius* mutant NGA23 were successful in that the same transformation frequencies (50-100 transformants per µg) as compared to the original procedure described above were obtained.

Klebe et al. (1983) describes that the PEG used is critical for reproducible transformation. Therefore Klebe et al. (1983) performed a time consuming PEG 1000 preparation for optimal transformation efficiency.

To circumvent this tedious procedure, the influence of several PEGs from Serva and Roth company was tested. Until now, the optimal transformation efficiencies were obtained using PEG 1000 from Roth company. Substitution of 1 M sorbitol in SBEG by 1.25 M KCl, 30 mM CaCl₂ leads to an increase in the transformation frequency by a factor 2, yielding 200 NGA23 transformants per µg YRpJD2 plasmid DNA.

During freezing of yeast cells, the plasma membrane becomes damaged, resulting in a higher permeability; in addition, the degree of cell damage is dependent on the velocity of freezing (Lepock et al., 1984). Based on these observations, the transformation competence of the cells was tested after freezing at -70°C in a freezer (relatively slow freezing), with dry ice acetone at -78°C (fast freezing) and with liquid nitrogen -196°C (very fast freezing).

Using liquid nitrogen, the transformation frequency decreased by 40%, whereas the usage of dry ice acetone increased the yield of transformants by 30% compared to freezing in a -70°C freezer.

The DNA was added to the frozen cells (stored at -70°C for 10 min up to 6 months) and thawed by rapid agitation at 37°C. Instead of rapid agitation in a 37°C water bath, an Eppendorf mixer, type 5432, at 37°C was used. This condition for rapid agitation during thawing gave the best reproducible transformation frequencies.

With these modifications, a transformation frequency of 300-1000 transformants per µg YRpJD2 plasmid DNA was obtained for *Schwanniomyces alluvius* NGA23 mutants.

With this simplified transformation procedure, high transformation frequencies could also be obtained for *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, and *Kluyveromyces lactis* (Table II).

Estimation of plasmid copy number in *Schwanniomyces alluvius* and *Saccharomyces cerevisiae* transformants

With our simplified transformation procedure, *Schwanniomyces alluvius* NGA23 was transformed with: pYe(trp5)1-53 - containing pBR313 (Bolivar et al., 1977) and the TRP5 gene of *Saccharomyces cerevisiae* pYAS1 - a 2 µM derivative containing both the LEU2 and TRP5 genes of *Saccharomyces cerevisiae* (Strasser, 1982)

YRP7TRP5 - containing ARS1 and both the TRP1 and TRP5 genes of *Saccharomyces cerevisiae*
YRpJD1 - containing both SwARS1 and ARS1 and both the TRP1 and TRP5 genes of *Saccharomyces cerevisiae*

YRpJD2 - containing SwARS1 and the TRP5 gene of *Saccharomyces cerevisiae*

pEK2TRP5 - containing both KARS1 and ARS1 and the TRP5 gene of *Saccharomyces cerevisiae*
The plasmids are summarised in Figure 8.

The transformation frequencies and stabilities of these plasmids in *Schwanniomyces alluvius* NGA23 are summarised in Table III. The copy numbers of the different plasmids were estimated by Southern blot analysis (Fig. 9 and 10).

The chromosomal HIS4 locus representing one copy per cell was used as an internal standard. There are three chromosomal EcoRI fragments (0.9 kb, 6.0 kb, 8.0 kb) which hybridize with ³²P-labelled YRP7HIS4 DNA. The homology of the chromosomal HIS4 locus and of the YRpJD1 plasmid to the hybridisation probe YRP7HIS4 is represented in Figure 10D.

For copy number estimation of the plasmids pYAS1, YRp7Trp4, YRpJD1, (YRpJD2) and pEK2TRP5, the intensity of the common 5.3 kb EcoRI band was compared with the 6 kb EcoRI band of the chromosomal HIS4 gene.

The intensity of staining in the autoradiogram reveals that the plasmids YRpJD1 and YRpJD2 have a copy number of about 5-10 copies per cell (Fig. 9 and 10).

Both the plasmids YRp7TRP5 and pEK2TRP5 have about the same average copy number per cell, which is in the range of about 0.2-1, independent of the orientation of the TRP5 fragment, whereas pYAS1 exists in about 2-3 copies per cell (Fig. 10).

ARS1 and 2 μ m DNA sequences of *Saccharomyces cerevisiae* can be used for autonomous replication in *Schwanniomyces alluvius* NGA23 although not as efficiently as SwARS1. SwARS1 behaves in *Schwanniomyces alluvius* as ARS1 in *Saccharomyces cerevisiae*, yielding 5-10 copies per cell (Fig. 11, Table IV). The plasmid YRpJD1 which has the ARS1 element in addition to SwARS1 element, showed no increased copy number as compared to YRpJD2, which only has the SwARS1 element. The plasmid pEK2TRP5, which has the KARS2 element in addition to the ARS1 element, has the same copy number as the plasmid YRp7TRP5, which only has the ARS1 element. It is therefore assumed that KARS2 cannot serve as an efficient autonomous replication element in *Schwanniomyces alluvius*.

The transformation of *Schwanniomyces alluvius* NGA23 with pYe(trp5)1-53 (Walz et al., 1978) can be explained by an abortive transformation event concerning the high instability of these transformants (Table III).

Regulation of the cloned *Schwanniomyces castellii* α -amylase gene and over-production of the protein in *Schwanniomyces alluvius* transformants

The production of α -amylase in the yeast species *Schwanniomyces* underlies catabolite repression in the presence of glucose (Sills et al., 1982).

The synthesis of α -amylase is induced by soluble starch dextrins and maltose (Sills et al., 1982). Maltose, which is part of the products after starch degradation, was found to be a stronger inducer than soluble starch (Sills et al., 1984b). It appears that a very low basal level of amylase is constitutively produced. This low level of α -amylase can hydrolyse starch, releasing low molecular weight sugar such as maltose, which can penetrate the cell membrane and in turn induce the synthesis of α -amylase (Sills et al., 1984b).

To study regulation and expression of the cloned α -amylase gene in *Schwanniomyces alluvius*, the 5 kb EcoRI fragment containing the structural α -amylase gene was inserted into the YRpJD2 (Fig. 8), resulting in plasmid YRpJD2- α and transformed into *Schwanniomyces alluvius* NGA23 with selection for tryptophan prototrophy on YNB plates.

As it is shown in Figure 5B, the total α -amylase activity of NGA23: YRpJD2- α transformants growing under induced condition in YNB maltose medium is increased 4-5 fold compared to control transformants NGA23: YRpJD2 which do not contain additional plasmid encoded α -amylase (Fig. 4A).

During the induction phase of α -amylase synthesis, the drastical 7-fold increase in extracellular α -amylase activity per cell concentration (OD_{600nm}) compared to control transformants, correlates with the estimated 5-10 copies of SwARS1 plasmids in *Schwanniomyces alluvius*.

It is therefore concluded that the increase in α -amylase activity is due to a gene dosage effect.

When the NGA23:YRpJD1 and NGA23:YRpJD2 transformants were grown under repressed conditions in YNB containing 2% glucose, no α -amylase activity could be measured in the extracellular medium until the glucose concentration dropped to about 0.2 mM (Fig. 4C and E). Thus the plasmid encoded α -amylase is regulated by catabolite repression as chromosomal α -amylase.

By construction of the α -amylase structural gene under control of a high expression promoter as e.g. that of PDC1 or PGK genes, higher expression strains independent of growth stage and C-source can be obtained.

The overproduction of the α -amylase protein was shown by analysis of *in vitro* labelled proteins of the culture supernatant by fluorography (Fig. 12). α -amylase activity could be measured in the extracellular medium of MC34:pYC1 α transformants under growth in YNB containing 2% and also 4% glucose as a sole carbon source indicating that the plasmid encoded α -amylase is not regulated by catabolite repression in *Saccharomyces cerevisiae*.

Isolation of auxotrophic mutants of *Schwanniomyces alluvius*

For prosecuting the cloning of the glucoamylase gene, the functional complementation of a glucoamylase mutant of *Schwanniomyces alluvius* was chosen.

A prerequisite is the availability of an appropriate mutant which can be complemented by the cloned gene. For this purpose, a *Schwanniomyces alluvius* mutant NGS1 which is not able to grow on maltose or soluble starch as a sole carbon source was analysed. When this mutant was grown on plates containing 1% glucose and 1% soluble starch, halo formation could be observed after iodine staining, indicating that α -amylase can be secreted by the mutant. No glucoamylase activity could be detected in crude extracts and culture supernatant.

As already mentioned, α -amylase can hydrolyse starch into dextrin, maltose and higher oligosaccharides (Fig. 7), whereas glucoamylase can use these products as substrates for the enzymatic hydrolysis into glucose. The interesting feature of the NGS1 mutant is the inability to grow on maltose. The *Schwanniomyces alluvius* wild type secretes glucoamylase which hydrolyses maltose to glucose in the extracellular medium, and hence, in contra to *Saccharomyces cerevisiae*, has no need for a maltose uptake system. It is therefore concluded that NGS1 is an α -amylase⁺ glucoamylase⁻ mutant, in having no functional glucoamylase either by a point mutation in the structural gene, or in the promoter region.

We have focused our attention on this glucoamylase mutant to clone the glucoamylase gene of *Schwanniomyces castellii* by functional complementation.

For cloning purposes, an additional mutation had to be introduced into the α -amylase⁺ glucoamylase⁻ mutant, preferentially a trp5 mutation (leading to a defect in the indole tryptophan reaction), because *Schwanniomyces alluvius* could be efficiently transformed using the TRP5 gene of *Saccharomyces cerevisiae* as a selectable marker on SwARS1 containing plasmids e.g. YRpJD1 and YRpJD2. In addition, these mutants can easily be identified by their property of indole accumulation.

For the isolation of trp5 auxotrophic mutants, all procedures were carried out at 30°C. 10 ml NGS1 culture was grown to OD₆₀₀ = 0.6 in complex medium, washed twice with H₂O and exposed to UV light (254 nm) for 50 seconds (1% survivals). The mutagenised cells were then incubated for 2 days on complex medium for segregation of the trp auxotrophs and plated on YEPD plates for replica plating on minimal medium containing tryptophan and on minimal medium plates containing all amino acids but no tryptophan.

From 2614 NGS1 colonies tested, 7 trp auxotrophic mutants which grow on minimal medium containing tryptophan but not on minimal medium containing all amino acids except tryptophan, could be isolated (reversion rate 10⁻⁷ - 10⁻⁸). Two of these seven mutants could be analysed to be defective in the indole tryptophan reaction since both were accumulating indole.

These trp5, α -amylase⁺glucose⁻ mutants were designated NJD1 and NJD2. These two strains could be efficiently transformed with YRpJD1 yielding approximately 800 transformants per μ g DNA, indicating that NJD1 and NJD2 are really mutated in the TRP5 gene.

Further cloning procedures can be carried out in an analogous manner as with the α -amylase gene.

Construction of an *E. coli* - *Saccharomyces cerevisiae* - *Schwanniomyces alluvius* cosmid shuttle vector for cloning genes in *Schwanniomyces alluvius*

In analogy to the cosmid pYc1 (2 μ) and pYc2 (ARS) which were constructed for cloning genes in *Saccharomyces cerevisiae* (Hohn and Hinnen, 1980), a cosmid for cloning in *Schwanniomyces alluvius* was constructed.

Cosmids are plasmids which contain the cos (cohesive sequence) region of bacteriophage lambda (Collins and Hohn, 1978) as well as the essential sequences for cloning (replication origins, a gene as selectable marker and suitable cloning sites). The cos sequence comprises the structural information necessary for packaging the DNA to produce an intact bacteriophage lambda (Hohn, 1975). High molecular oligomeric cosmid DNA can therefore be efficiently packaged by an *in vitro* packaging system (Hohn, 1979). If the cos sequences are separated by any 37-52-kb DNA fragments corresponding to about 73% to 105% of the lambda phage genome (Feiss et al., 1977). The relatively large size of clonable fragments presents the possibility that only about 1,200 cosmid clones containing fragments with an average size of about 30-40 kb are necessary to encompass the genome of *Saccharomyces cerevisiae* by more than 95% (Hohn and Hinnen, 1980).

For cloning large DNA fragments, the advantage in using cosmid instead of plasmid DNA is that relatively high molecular oligomeric DNA can easily be obtained by ligation of highly concentrated DNA, whereas the efficiency to form circular transformable plasmids by ligation of large DNA fragments with smaller vector DNA is disadvantageous because of the relation between the fragment size and effective concentration of free ends (Dugaiczky et al., 1975). In addition, the packaged DNA can efficiently be transduced by infection of lambda sensitive *E. coli* cells (Hohn and Hinnen, 1980).

Based on these observations, an *E. coli* - *Schwanniomyces alluvius* - *Saccharomyces cerevisiae* cosmid shuttle vector was constructed by subcloning the 1.7 kb BglII fragment of plasmid pHC79 containing the cos region (Hohn and Collins, 1981) into one of the two BamHI sites of YRpJD2. YRpJD2 was therefore partially digested with BamHI (0.5 units/ μ g DNA at 30°C) and the resulting largest linear DNA which arises from a single cleavage by BamHI was isolated by preparative gel electrophoresis and ligated with the 1.7 kb BglII fragment of pHC79, which has also been isolated by preparative gel electrophoresis after complete digestion of pHC79 DNA. This ligation mixture was used to transform *E. coli* JA221, selecting on ampicillin resistance. Subclones were screened by colony hybridisation using the nick translated 1.7 kb BglII fragment as a radioactive probe.

From the positive clones, two different cosmids, namely pCJD5-1 and pCJD5-2 (Fig. 13a) could be isolated and analysed after cleavage with EcoRI, BamHI and PvuII restriction enzymes (Fig. 13b). Figure 13b demonstrates that in cosmid pCJD5-1, the cos sequence is subcloned into the BamHI site of YRpJD2 between the SwARS1 sequence and the TRP5 gene. In pCJD5-2, the cos region was inserted in tandem at the same position. The existence of two, in tandem, orientated cos elements should drastically increase the packaging efficiency compared to cosmids which contain only a single cos element (Lindenmeier et al., 1982). Up till now, it was not tested whether this is also the case for pCJD5-2.

With cosmids pCJD5-1 and pCJD5-2, 500-1000 NJD1 transformants per μ g cosmids DNA were obtained by selecting on tryptophan, indicating that the cos element does not influence the transformation frequency. Both cosmids contain

- (a) the SwARS1 sequence which acts as replication origin in *Schwanniomyces alluvius* as well as *Saccharomyces cerevisiae*
- (b) the origin for replication of *E. coli*;
- (c) the ampicillin resistance gene as selectable marker in *E. coli*;
- (d) the TRP5 gene of *Saccharomyces cerevisiae* as selectable marker in *Schwanniomyces alluvius* as well as in *Saccharomyces cerevisiae* trp5 mutants; and
- (e) the cos region as prerequisite for the construction of a cosmid library.

In addition, both cosmids contain suitable cloning sites, namely BamHI and Sall for cloning Sau3A, BamHI, BglII, Sall and XHOI fragments, and hence provide a new class of cosmid vectors for the construction of a genomic DNA library to clone genes in *E. coli*, *Saccharomyces cerevisiae* and *Schwanniomyces alluvius*.

Cloning of the glucoamylase gene from *Schwanniomyces castellii*

The cloning of the glucoamylase gene was started from a *Schwanniomyces castellii* ATCC 26076 cosmid pool which was established by ligating partially BamHI digested genomic DNA of *Schwanniomyces castellii* into the *E. coli* - *Saccharomyces cerevisiae* - *Schwanniomyces alluvius* shuttle vector pCJD5-1 using the cosmid cloning technique described by Ish-Horowitz et al., 1981.

After *in vitro* packaging of the DNA ligation mixture, *E. coli* strain HB101 was infected, yielding approximately 15,000 ampicillin resistant colonies corresponding to approximately 30 genome size equivalents of *Schwanniomyces castellii* assuming the same genome size for *Schwanniomyces castellii* as for *Saccharomyces cerevisiae* (approx. 15,000 kb) and a cloned DNA fragment ranging between 30 and 40 kb in size. From these transductants, cosmid DNA was isolated and used for the transformation of NJD1 (α -amylase*glucoamylase*trp5) with selection for trp prototrophy on minimal medium containing 2% glucose.

The transformation frequencies using both the protoplast method (Beggs, 1978) and the simplified freeze thaw method was relatively low (1-10 transformants/ μ g cosmid DNA). Nevertheless approx. 3,000 transformants could be obtained which were screened for their regained ability to grow on soluble starch as sole carbon source. One NJD1 transformant which has a Trp⁺ and glucoamylase⁺ phenotype could be isolated from starch plates. In a mitotic stability test (Beggs, 1978), both the TRP5 phenotype and the glucoamylase complementing functions were segregated out, simultaneously indicating that both genes were carried by the cosmid. The cosmid was rescued by transforming *E. coli* HB101 with yeast minilysate

and selection for ampicillin resistance. Cosmid DNA was isolated and used for retransformation of NJD1, to ascertain that the glucoamylase complementing function is still carried by the isolated cosmid DNA. After cleavage of the cosmid with BamHI, two BamHI sites could be localised within the cloned fragment, dividing it into the three fragments, two of them being of approx. 10 kb and 12 kb.

In order to decrease the size of the approx. 40 kb size cosmid, BamHI fragments were subcloned into the single BamHI site of pCJD5-1 plasmid. Therefore, the cosmid DNA was digested with BamHI, religated and transformed into *E. coli* HB101. The plasmid DNA from ampicillin resistant cells was used to transform NJD1 and to screen for growth on soluble starch as a sole carbon source. Plasmid DNA was rescued by transforming HB101 with yeast minitylate and selection for ampicillin resistance. Plasmid DNA was isolated for restriction analysis after digestion with BamHI endonuclease.

Figure 15 demonstrates that the glucoamylase complementing function is carried by the 12 kb BamHI fragment, subcloned into pCJD5-1. This plasmid is designated pCJD5-AMG1 (Fig. 14). Since this plasmid contains the SwARS1 element acting in *Saccharomyces cerevisiae* as well as in *Schwanniomyces alluvius*, the cloned gene in *Saccharomyces cerevisiae* was analysed by transforming X3656 selecting on tryptophan prototrophy. Unfortunately the transformants were not able to use soluble starch as a sole carbon source. The first explanation, that the 5-10 copies of pCJD5-AMG1 in *Saccharomyces cerevisiae* are not high enough to overcome possible proteolytic degradation, failed, since after subcloning of the approx. 12 kb BamHI fragment into the high copy number, plasmid pJDB207 (Beggs, 1981) (Fig. 14) having 50-100 copies per cell, and transformation of GRF18 *Saccharomyces cerevisiae* cells, none of the transformants were able to grow on soluble starch as a sole carbon source.

Although glucoamylase activity could be measured in NJD1:pCJD5-AMG2 transformants, no activity in both crude extract and culture filtrate of *Saccharomyces cerevisiae* transformants could be detected. Because Käufer et al. (1985) have recently shown that *Schizosaccharomyces pombe* has a considerable advantage over *Saccharomyces cerevisiae* in accurately excising an intervening sequence from a transcript of a higher eucaryotic gene, attempts were made to express the glucoamylase in this fission yeast. However, no expression and secretion of glucoamylase could be detected in *Schizosaccharomyces pombe* leu1-32 (complemented with LEU2 of *Saccharomyces cerevisiae* after transformation with plasmid pJDB207-AMG1 (Fig. 14). The transformation of the gene into *Kluyveromyces lactis* SD11(trp1) with pEK2-AMG2 (Fig. 14) did not lead to transformants which can grow on soluble starch as sole carbon source. It is therefore concluded that a *Schwanniomyces* gene was cloned which cannot be functionally expressed in *Saccharomyces cerevisiae*, *Schizosaccharomyces pombe*, and *Kluyveromyces lactis* because of different splicing specificities, promoter recognition of *Schwanniomyces* specific post transcriptional processing.

For further analysis, subcloning smaller fragments of the 12 kb DNA was carried out in order to find the smallest possible DNA fragment still carrying the functional complementation property. The molecular weight of glucoamylase is approx. 117,000 d (Simoes-Mendes, 1984) corresponding to a long structural gene of about 3 kb in length.

Furthermore, the gene can be sequenced to analyse codon usage and possibly existing introns. Additionally, the 5' non-coding leader sequence of *Schwanniomyces castellii* DNA can be exchanged by a well characterised promoter such as the ADH1 (Hitzeman et al., 1981) or PDC promoter (Hollenberg et al., 1983) to exclude possible inefficiency of the *Schwanniomyces* promoter in other yeast species.

In order to characterize the gene, glucoamylase was isolated for the production of glucoamylase specific antibodies to assay the crude extracts and culture filtrate of NJD1 as well as *Saccharomyces cerevisiae* or *Schizosaccharomyces pombe* and *Kluyveromyces lactis* transformants for the presence of inactive glucoamylase. No promoter activity could be detected in *Saccharomyces cerevisiae* transformants by Northern analysis. In addition there was no protein detectable in culture supernatants and crude extracts of the *Saccharomyces cerevisiae* transformants using a specific antibody raised against glucoamylase. In addition, peptides of purified glucoamylase were isolated for amino acid sequencing in order to synthesize synthetic oligonucleotides for analysing the cloned gene.

To exclude cloning artefacts arising from the construction of the *Schwanniomyces* cosmid library a Charon 4A *Schwanniomyces castellii* library, as described by Maniatis et al. 1982, containing partial EcoRI fragments (10-30kb) of *Schwanniomyces castellii* DNA yielding approximate 150000 inserts was screened directly for the glucoamylase gene.

The Charon library was further analysed by two different oligonucleotide mixtures which were deduced from a part of peptide 3 (amino acid 11-18) and from peptide 4 respectively (see Fig. 3). The smallest restriction fragment hybridizing with both 5' labelled oligonucleotide mixtures was a 3.7kb BglII DNA fragment. S1 mapping revealed that a complete transcript maps within that fragment. This fragment was sequenced by Maxam and Gilbert (1980). The DNA sequence is presented in Fig. 17.

All five peptide sequences, (Fig. 3) were found within the open reading frame of 2874 bases.

The expression of the glucoamylase gene can be achieved by promoter exchange. A Saccharomyces cerevisiae promoter region initiation of transcription and regulation when inserted 5' upstream to the coding sequence of the glucoamylase gene. There are no suitable restriction sites in front of the translation start point of the glucoamylase gene which could be used for promoter fusions. Therefore, we introduced a BamHI and SalI restriction site 5' to the translation initiation codon by site-directed mutagenesis by the gapped duplex DNA method (Kramer et al (1984)).

A synthetic oligonucleotide of 42 bases (5' CTCATGACTGTGTCGACGGATCCAAGATGATTTTCT-GAAGC 3') was annealed to the gapped duplex DNA. Filling up the single stranded areas with DNA polymerase (large fragment), ligation and E. coli transformation were performed and a recombinant 10 mutagenized plasmid could be isolated containing a BamHI site at position -9 and a Sall site at position -15 with respect to the first base of the translation initiation codon.

From the resultant recombinant plasmid the structural gene can be sub-cloned as a BamHI/SalI, BamHI/HindIII, SalI/SalI, BamHI/PstI, SalI/PstI, BamHI/XbaI and SalI/XbaI in appropriate expression vectors fusing the gene with one of the mentioned restriction sites to different promoters. Every other suitable restriction site can be introduced by a suitable linker.

Different expression vectors using different promoters, different selectable markers and different replication origins (2 μ , ARS and CEN) for yeast were constructed.

For example, the gene was fused to the inducible galactokinase (GAL1) promoter in plasmid pBM272, which is directly derived from vector pBM150 (Johnston and Davis 1984) by introducing a Hind III site immediately behind the BamHI site. The DNA sequence of the fusion is as follows:

5' upstream activation site
GATATATAAATGCAAAACTGCATAACCACTTTAACTAATACTTTCAACATTTTCGGTTTGTATTACTTCT-
 TATTCAAATGTAATAAAAGTATCAACAAAAAATTGTTAATATACCTCTATACTTTAACGTCAAGGAGAAAAAA-
 CCCCCGATCCAAGATGATTTTTCTGAAGC....3'

25 The fragment containing the glucoamylase gene was sub-cloned into the vector pBM272 as a BamHI/Sall fragment. The resulting plasmid was transformed into Saccharomyces cerevisiae α MZ4 (α gal1, trp1, ura3, leu2). This leads to inducible expression of the glucoamylase gene and to secretion of active glucoamylase.

The amount of activity in the culture supernatant was in the range which was measured in culture supernatants of *Schwanniomyces alluvius* grown under induced conditions.

The glucoamylase secreted by the *Saccharomyces cerevisiae* transformant has maximum activity at 50°C and can be inactivated at 60°C as is the case for *Schwanniomyces alluvius* glucoamylase.

The expression of the glucoamylase and secretion of the glucoamylase protein could also be achieved under control of the pyruvate decarboxylase (PDC) promoter (Hollenberg et al., 1983), phosphoglycerate kinase (PGK) promoter (Dobson et al., 1982), glyceraldehyde-3-phosphate dehydrogenase (GAPDH) promoter of the gene on pgap491 (Holland and Holland, 1979), alcoholdehydrogenase 1 (ADH1) promoter (Hitzeman et al., 1981), copper-chelatin (CUP1) promoter (Karin et al., 1984, Butt et al., 1984) and alcoholdehydrogenase 2 (ADH2 or ADH2) promoter (Russell et al., 1983, Beijer et al., 1985).

Over production of glucoamylase could be achieved using the PDC-promoter and high copy number plasmid (2 μ with leu2d). Expression of the glucoamylase gene and secretion of the glucoamylase protein can also be detected in Kluyveromyces lactis after fusion of the transcriptional signals of the β -galactosidase gene (Breunig et al., 1984).

Expression of the glucoamylase gene and secretion of the glucoamylase protein can also be detected in Schizosaccharomyces pombe after fusion of the transcriptional signals of the alcohol dehydrogenase (ADH1) gene (Russell et al., 1983).

Expression of the glucoamylase gene and secretion of the glucoamylase protein can also be detected in Hansenula polymorpha after fusion of the transcriptional signals of dihydroxyacetone synthase (DHAS) (Janowicz et al., 1985) methanol oxidase (MOX) (Ledeboer et al., 1985).

In addition secretion of the α amylase could also be detected after exchange of the Schwanniomyces
50 promoter by the MOX promoter in Hansenula polymorpha.

The glucoamylase gene was integrated into the chromosom of Saccharomyces cerevisiae without E. coli sequences. Therefore, the different promoter-glucoamylase fusions were inserted into a suitable DNA sequence of Saccharomyces cerevisiae, preferably the homothalium gene (HO) or an ARS-sequence. A linear DNA fragment could then be generated which has ends within the Saccharomyces cerevisiae sequences flanking the functional glucoamylase gene. By cotransformation of Saccharomyces cerevisiae with YEP 36 (Butt et al., 1984) the gene could be stably integrated. The integrants exhibit glucoamylase activity in the culture supernatant and has the same fermentation and growth properties as the wildtype.

Genetically manipulated microorganisms, preferably yeasts, as described above in this specification are of great advantage in a lot of fermenting processes because of their ability to hydrolyse higher molecular carbohydrates.

One of the possible uses of the genetically altered yeast strains described above is in the production of biomasses. Since the yeast strains having the new ability to secrete α -amylase and glucoamylase still remained there other good fermentation abilities biomasses can be produced most efficiently by the use of the inventive yeast strains. The methods of producing the biomasses are the usual ones, known by the man skilled in the art.

A further field of use of Saccharomyces cerevisiae, having the new ability of producing α -amylase and glucoamylase is the production of bakers yeast and of bread, using the bakers yeast.

The inventive yeast can be used as pellet yeast, for example, made by conventional drum-drying processes or can be powdered yeast, for example, having a particle size of 1.7 mm or less. Making the powdered yeast comprises spray-drying of a liquid yeast composition in air, followed by further drying, as known in the state of the art. The inventive yeast, for example, may be prepared as a dried yeast as described in GB-PS 1 459 407, GB-PS 1 459 085 or GB-PS 1 230 205. Any other methods known in the art for producing bakers yeast or producing baked products by using the new yeast of this invention are also suitable.

The genetically manipulated yeast strains provided according to this invention are further highly suitable for the production of ethanol. The preferred organism for the production of ethanol by fermentation is the yeast Saccharomyces cerevisiae and the preferred carbohydrate used in the ethanol production is a starch. Thus, a Saccharomyces cerevisiae, being able to ferment starch quantitatively is highly advantageous in the production of ethanol. The production of potable spirit or industrial ethanol by the use of the genetically manipulated yeast strain of the present invention can be carried out in a manner known per se. The inventive yeast strain has the ability to ferment concentrated carbohydrate solutions, has ethanol tolerance and the ability to produce elevated concentrations of ethanol, has a high cell viability for repeated recycling and has temperature tolerance.

Possibly the most important field of use of the new Saccharomyces cerevisiae, being able to produce α -amylase and glucoamylase is the production of special beers, preferably low carbohydrate beers. As already described in the specification above, one technique for producing low carbohydrate beers was to add amylolytic enzymes to the wort during fermentation. Because 70% to 75% of the dextrins in wort are of the branched type, debranching enzyme is essential for total hydrolyses of wort dextrins to fermentable sugars. The glucoamylase, produced by the new genetically manipulated Saccharomyces cerevisiae possesses debranching activity and therefore can hydrolyse the dextrins completely. An important characteristic of the amylolytic enzymes produced by the new yeast Saccharomyces cerevisiae is their sensitivity to the normal pasteurisation cycle employed in brewing.

The method of preparing beers has a series of stages, starting with malting, where barley is soaked in water and allowed to germinate and during this step amylase is released which converts the starch from the grain into sugar. During the next step, grains are removed by filtration and the resulting liquid wort is brewed with hops. The hops give the beer the bitter flavour and during this step the enzymatic conversion of starch to sugar is stopped. Next, the wort is cooled and filtered to remove the precipitated proteins. Then yeast is added to the wort to convert the sugars to alcohol and carbon dioxide in a fermentation process, after which the mixture is stored, the yeast is settling out, the beer is filtered again, bottled or canned and pasteurised.

In order to produce a low-calorie beer, it is desirable to consume as many of the carbohydrates as possible in the fermentation process. For this purpose the new Saccharomyces cerevisiae, being able to produce α -amylase and glucoamylase is highly suitable.

A method for preparing the low-calorie beer can be carried out generally as described in the European Patent Application 0 163 135, with the exception that no amylolytic enzymes have to be added during the brewing process. The above-described usual brewing steps can be carried out without any time-consuming or expensive additions of amylolytic enzymes. Carrying out the brewing process as known in the art, using the new genetically manipulated Saccharomyces cerevisiae according to this invention, results in a special beer, preferably low-carbohydrate beer, low-calorie beer depending on the degree of the fermentation of sugar. The brewing process, using the Saccharomyces cerevisiae of this invention is decreased in time due to a more rapid decrease in E_s at the end of the fermentation process.

Using the Saccharomyces cerevisiae of this invention, the manufacturing process can be adapted for the production of a beer from a high gravity wort, the carbohydrate level of which can be regulated by a predetermined fermentation time and dilution thereafter. Furthermore, the manufacturing process may be adapted for the fermentation of substrates, the constituent of which are for a considerable part suspended hydrated carbohydrates. The genetically engineered Saccharomyces cerevisiae can be brought into contact with the fermented wort subsequently to main fermentation but prior to conditioning.

A more detailed description of the production of a low-calorie beer is now given in the following examples.

10

Example I

Brewers' wort was prepared using conventional brewing techniques in such a way that 80% of the extract in the wort was derived from brewer's malt and 20% derived from corn grits. This wort had an original gravity of 11.7 °P. 3 litre fleakers were filled with 2.5 litres of aerated wort and pitched with 1.5 g/l wet yeast of a genetically manipulated Saccharomyces cerevisiae lager brewing strain, which had received by transformation the α -amylase and/or glucoamylase gene (s) from Schwanniomyces castellii. Both enzymes were secreted by the transformed yeast strains. As a control the original production lager strain of Saccharomyces cerevisiae was used.

Fermentation was allowed to proceed at a controlled temperature of 9°C under anaerobic conditions and with mechanical stirring due to the small volume of fermentation. At the end of the fermentation, the beer was cooled to 0°C, yeast was separated from the beer and analysed for residual extract and % ethanol. Also a typical gas chromatogram of the green beer was made to determine the amount of esters and higher alcohols.

Analysis of the green beer fermented with the original lager brewing strain (A), the genetically manipulated brewing lager strain, secreting α -amylase (B) or glucoamylase (C) or both enzymes α -amylase and glucoamylase (D).

30

35

40

45

50

55

		A	B	C	D	
5	original gravity ^o P	11.70	11.53	11.60	11.57	
	ethanol % w/w	3.99	4.02	4.72	4.74	
10	apparent extract ^o P	2.14	1.88	0.30	0.22	
	apparent degree of fermentation %	81.7	83.7	97.4	98.1	
	real extract ^o P	3.96	3.71	2.43	2.36	
15	real degree of fermentation %	66.2	67.8	79.0	79.6	
20	Gas chromatografic analysis					
	Acetaldehyde	mg/l	2.0	2.1	3.1	3.9
25	Acetone	mg/l	0.29	0.37	0.38	0.38
	Ethyl formiate	mg/l	0.05	0.18	0.19	0.14
	Ethyl acetate	mg/l	20.6	21.7	33.0	39.9
30	Ethyl propionate	mg/l	0.07	0.06	0.06	0.07
	Isoamyl acetate	mg/l	2.45	2.21	2.95	2.92
	Methanol	mg/l	2.0	2.5	2.3	2.6
	N-propanol	mg/l	17.5	16.3	21.0	17.7
35	Isobutanol	mg/l	14.2	12.6	14.7	15.3
	Opt.act.amylalcohol	mg/l	21.5	19.0	21.8	20.4
	Isoamyl alcohol	mg/l	62.5	54.2	57.6	55.1
40	Total higher alcohols	mg/l	115.7	102.1	115.1	108.5

Example II

The fermentation process was the same as for Example I except that the fermentation temperature was 13.5°C. After 7 days of fermentation the beer was cooled to 0°C.

Analysis of the green beer fermented with the original lager brewing strain (A), the genetically manipulated brewing lager strain, secreting α -amylase (B) or glucoamylase (C).

50

55

		A	C
5	original gravity °P	11.8	11.8
	ethanol % w/w	4.10	4.60
	apparent extract °P	2.00	0.85
10	apparent degree of fermentation %	83.1	92.8
	real extract °P	3.90	2.90
15	real degree of fermentation %	67.0	75.4

The analysis of the beers fermented with a genetically manipulated lager brewing strain, secreting α -amylase (B) was hardly different from the control.

Example III

The fermentation process was the same as for Example I except that the prepared wort had an original gravity of 15°P. After 9 days of fermentation the beer was cooled to 0°C.

Analysis of the green beer fermented with the original lager brewing strain (A), the genetically manipulated brewing lager strain, secreting α -amylase (B) or glucoamylase (C).

		A	C
30	original gravity °P	15.0	15.0
	ethanol % w/w	4.57	5.22
35	apparent extract °P	4.30	2.76
	apparent degree of fermentation %	71.4	81.6
40	real extract °P	6.31	5.06
	real degree of fermentation %	58.0	66.3

The analysis of the beers fermented with a genetically manipulated lager brewing strain, secreting α -amylase (B) was hardly different from the control.

Example IV

The fermentation process was the same as for Example I except that the prepared wort had an original gravity of 8.23°P and the fermentation time/temperature profile was as follows:
4 days 9.5°C, 3days 13.5°C.

Analysis of the green beer fermented with the original lager brewing strain (A), the genetically manipulated brewing lager strain, secreting α -amylase (B) or glucoamylase (C).

		A	C
5	original gravity $^{\circ}\text{P}$	8.23	8.23
	ethanol % w/w	2.88	3.60
	apparent extract $^{\circ}\text{P}$	1.16	-0.56
10	apparent degree of fermentation %	85.9	107.0
	real extract $^{\circ}\text{P}$	2.48	1.14
15	real degree of fermentation %	69.9	86.7

The analysis of the beers fermented with a genetically manipulated lager brewing strain, secreting α -amylase (B) was hardly different from the control.

20

Example V

The fermentation process was the same as for Example I except that the prepared wort consisted of 66.7% standard brewers' wort as described in Example I and 33.3% suspended hydrated starch (12% w/v).

25 Analysis of the green beer fermented with the original lager brewing strain (A), the genetically manipulated brewing lager strain, secreting α -amylase (B) or glucoamylase (C) or both enzymes α -amylase and glucoamylase (D).

30

35

40

45

50

55

		A	B	C	D
5	original gravity ^o P	12.01	12.14	12.13	12.22
	ethanol % w/w	2.86	4.03	4.19	4.96
10	apparent extract ^o P	5.15	2.49	1.63	0.37
	apparent degree of fermentation %	57.1	79.5	86.6	96.9
15	real extract ^o P	6.47	4.32	3.62	2.61
	real degree of fermentation %	46.2	64.4	70.2	78.6
20	Gas chromatographic analysis				
	Acetaldehyde mg/l	3.9	3.4	8.4	11.6
25	Acetone mg/l	0.24	0.20	0.17	0.23
	Ethyl formiate mg/l	0.07	0.13	0.22	0.25
	Ethyl acetate mg/l	8.7	20.5	39.6	43.3
30	Ethyl propionate mg/l	-	0.02	0.01	0.01
	Isoamyl acetate mg/l	0.62	1.25	1.62	1.70
	Methanol mg/l	-	-	-	-
35		A	B	C	D
	N-propanol mg/l	13.6	14.9	18.3	19.7
	Isobutanol mg/l	13.0	11.5	14.7	16.5
40	Opt.act.amylalcohol mg/l	10.4	12.5	20.1	18.2
	Isoamyl alcohol mg/l	50.9	57.1	67.5	79.2
	Total higher alcohols mg/l	87.9	96.0	120.5	133.6

45
Example VI

Milled grains of wheat and maize were suspended in an appropriate buffer, boiled in order to gelatinaze the starch and liquified by treatment with a commercial α -amylase enzyme preparation.

50 This broth had an original gravity of 18.9^oP and was fermented in 2.5 litres fermentors at a temperature of 30°C for 72 hours. The yeast strain used was a Saccharomyces cerevisiae distilling strain, which had received by transformation the glucoamylase gene of Schwanniomyces castellii. Glucoamylase was secreted by the transformed yeast strain. As a control the same fermentation was carried out with the original untransformed distilling strain. After 72 hours the broth was filtered and the ethanol concentration was

55 determined.

Yeast	ethanol (% w/w)
untransformed distilling strain	8.9
transformed distilling strain	9.6.

REFERENCES

- BEIER, D.B., SLEDZIEWSKI, A., YOUNG, E.T. (1985) *Mol. Cell. Biol.* **5**, 1743-1749.
- BENOIST, C., O'HARA, K., BREATHNACH, R. and CHAMON, P. (1980) *Nuc. Acid. Res.* **8**, 127-142.
- 5 BEGGS, J.D. (1978): Transformation of yeast by a replicating hybrid plasmid. *Nature* **275**, 104-108.
- BEGGS, J.D. (1981): In *Molecular Genetics in Yeast*, Alfred Benzon Symposium **16**, 383-395.
- BIRNBOIM, H.C. and DOLY, S. (1979): A rapid alkaline extraction procedure for screening recombinant plasmid DNA
- 10 *Nuc. Acid. Res.* **7**, 1513-1523.
- BOLIVAR, S., RODRIGUEZ, R.L., BETLACH, M.C. and BOYER, H.W. (1977): Construction and characterisation of new cloning vehicles. I ampicillin-resistant derivatives of the plasmid pMB9. *Gene* **2**, 75-93.
- BRADFORD, M. (1976): A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein-dye binding.
- 15 *Anal. Biochem.* **77**, 248-254.
- BREUNIG, K.D., DAHLEMS, U., DAS S., and HOLLENBERG, C. P. (1984) *Nucl. Acids Res.* **12**, 2327-2341.
- BUTT, T.R., STERNBERG, E.J., GORMAN, J.A., CLARK, P., HAMER, D., ROSENBERG, M., and CROOKE, S.T. (1984) *Proc. Natl. Acad. Sci. USA* **81**, 3332-3336.
- 20 CARBON, J., RATZKIN, B., CLARKE, L. and RICHARDSON, D. (1977): The expression of cloned eucaryotic DNA in procaryotes. *Brookhaven Symp. Biol.* **29**, 277-296.
- COHEN, J., GORMAN, J., KOLTIN, Y. and DE WILDE, M. (1985): Molecular cloning of a glucoamylase gene of *Candida albicans*: Expression and secretion in *Saccharomyces cerevisiae*.
- 25 *Cold Spring Harbour Symp. Molec. Biol. of Yeasts*, **345**.
- COLLINS, J. and HOHN, B. (1978): Cosmids: A type of plasmid genecloning vector that is packageable in vitro in bacteriophage Lambda heads. *Proc. Natl. Acad. Sci. USA* **75**, 4242-4246.
- 30 DAS, S. and HOLLENBERG, C.P. (1982): A high-frequency transformation system for the yeast *Kluyveromyces fragilis*. *Curr. Genet.* **6**, 123-128.
- DAS, S., KELLERMANN, E. and HOLLENBERG, C.P. (1984): Transformation of *Kluyveromyces fragilis*. *J. Bacteriol.* **158**, 1165-1167.
- DAVIS, R.W., BOTSTEIN, D. and ROTH, J.F. (1980): Advanced bacterial genetics. A manual for genetic engineering. Cold Spring Harbour Laboratory.
- 35 DOBSON, M.J., TUIE, M.F., ROBERTS, N.A., KINGSMAN, A.J., KINGSMAN, S.M., PERKINS, R.E., CONROY, S.C., DUNBAR, B. and FOTHERGILL, L.A. (1982) *Nucleic Acids Research* **10**, 2625-2637.
- 40 DUGAICZYK, A., BOYER, H.W. and GOODMAN, H.M. (1975): Ligation of EcoRI endonuclease generated fragments into linear and circular structures. *J. Mol. Biol.* **96**, 171-184.
- FEISS, M., FISHER, R.A., CRAYTON, M.A. and EGNER, C. (1977): Packaging of the bacteriophage Lambda chromosome: Effect of chromosome length.
- 45 *Virology* **77**, 281-293.
- FERGUSON, A.R., KATSUNUMA, T., BETZ, H. and HOLZER, H. (1973): Purification and properties of an inhibitor of the tryptophan-synthetase-inactivating enzymes in yeast. *Eur. J. Biochem.* **32**, 444-450.
- FOGARTY, W.M. and KELLY, C.T. (1979): Starch degrading enzymes of microbial origin. In: BULL, M.J., (ed): *Progress in industrial microbiology*. Elsevier Scientific Publishing Company, New York,
- 50 **15**, 87-150.
- GAFNER, J., DE ROBERTIS, E.M. and PHILIPPSEN, P. (1983). *EMBO J.* **2**, 583-591.
- GANNON, F., O'HARA, K., PERRIN, F., LE PENNEC, J.P., BENOIST, C., COCHET, M., BREATHNACH, R., ROYAL, A., GARAPIN, A. and CHAMON, P. (1979): *Nature* **278**, 248-434.
- 55 HITZEMAN, R.A., HAGIE, F.E., LEVINE, H.L., GOEDEL, D.V., AMMERER, G. and HALL, B.D. (1981): Expression of a human gene for interferon in yeast. *Nature* **294**, 717-722.
- HOHN, B. (1975): DNA as substrate for packaging into bacteriophage lambda in vitro.

- J.Mol.Biol. 98, 93-106.
- HOHN, B. (1979): In vitro packaging of Lambda and cosmid DNA. In WU, R. (ed): Recombinant DNA, Methods In Enzymology. Academic Press, New York 68, 299-309.
- 5 HOHN, B. and COLLINS, J. (1980): A small cosmid for efficient cloning of large DNA-fragments. *Gene* 11, 291-298.
- HOHN, B. and HINNEN, A. (1980): Cloning with cosmids in E.coli and yeast. In SETLOW, J.K. and HOLLAENDER, A. (ed.): Genetic engineering - principles and methods. Plenum Press, New York & London 2, 169-183.
- 10 HOLLAND, J.P. and HOLLAND, M.J. (1979) *J. Biol. Chem.* 254, 5466-5474.
- HSIAO, C.L. and CARBON, J. (1981 b): Characterisation of a yeast replication origin (ars2) and construction of stable minichromosomes containing cloned centromere DNA (CEN3). *Gene* 15, 157-166.
- INNIS, M.A., HOLLAND, M.J., McCABE, P.C., COLE, G.E., WITTMAN, V.P., TAL, R., WATT, K.W.K.,
- 15 GELFAND, D.H., HOLLAND, J.P. and MEADE, J.H. (1985): Expression, glycosylation and secretion of an *Aspergillus glucoamylase* by *Saccharomyces cerevisiae*. *Science* 228, 21-26.
- ISH-HOROWICZ, D. and BURKE, J.F. (1981): Rapid and efficient cosmid vector cloning. *Nuc. Acids. Res.* 9, 2989-2998.
- 20 ITO, H., FUKUDA, Y., MURATA, K. and KIMURA, A. (1983): Transformation of intact yeast cells treated with alkali cations. *J.Bacteriol.* 153, 163-168.
- JANOWICZ, Z.A., ECKART, M.R., DREWKE, C., ROGGENKAMP, R.O., HOLLENBERG, C.P., MAAT, J., LEDEBOER, A.M., VISSER, C. and VERRIPS, C.T. (1985)
- 25 *Nucleic Acids Res.* 9, 3043-3062.
- JOHNSTON, M. and DAVIS, R.W. (1984) *Mol. Cell.Biol.* 4, 1440-1448.
- KARIN, M., NAJARIAN, R., HASLINGER, A., VALENZUELA, P., WELCH, J. and FOGEL, S. (1984) *Proc. Natl.Acad.Sci. USA* 81, 337-341.
- KATSUNUMA, T., SCHÖTT, E., ELSÄSSER, S. and HOLZER, M. (1972):
- 30 *Eur.J.Biochem.* 27, 520-526.
- KÄUFER, N.F., SIMANIS, V. and NURSE, P. (1985): Fission yeast *Schizosaccharomyces pombe* exises a mammalian RNA transcript intervening sequence. *Nature* 318, 78-80.
- KLEBE, R.J., HARRISS, J.V., SHARP, Z.D. and DOUGLAS, M.G. (1983): A general method for polyethylene-glycol-induced genetic transformation of bacteria and yeast. *Gene* 25, 333-341.
- 35 KRAMER, W., DRUTSA, V., JANSEN, H.-W., KRAMER, B., PFLUGFELDER, M. and FRITZ, H.-J. (1984) *Nucl.Acids.Res.* 12 9441-9456.
- LAEMMLI, U.K. (1970): Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 227, 680-685.
- 40 LEDEBOER, A.M., MAAT, J., VISSER, C., BOS, J.W., VERRIPS, C.T., JANOWICZ, Z., ECKART, M., DREWKE, C., ROGGENKAMP, R. and HOLLENBERG, C.P., (1985) *Nucl.Acids.Res.* 9, 3063-3082.
- LEPOCK, J.R., KEITH, A.D. and KRUUV, J. (1984): Permeability changes in yeast after freeze, thaw damage; comparison to reproductive survival.
- 45 *Cryo-Letts.* 5, 277-280.
- LINDENMAIER, W., HAUSER, H., GREISER DE WILKE, I., and SCHÜTZ, G., (1982): Gene shuttling; moving of cloned DNA into and out of eucaryotic cells. *Nucl. Acids. Res.* 10, 1243-1256.
- MANIATIS, T., FRITSCH, E.F. and SAMBROCK, J. (1982): Molecular cloning, a laboratory manual.
- 50 Cold Spring Harbour Laboratory, New York.
- MANNEY, T.R., DUNTZE, W., JANASKO, N. and SALAZAR, J. (1969): Genetic and biochemical studies of partially active tryptophane-synthetase mutants of *Saccharomyces cerevisiae*. *J. Bacteriol.* 99, 590-596.
- MAXAM, S.M. and GILBERT, W. (1980): Sequencing end labelled DNA with base specific chemical
- 55 cleavages. *Methods Enzymol.* 65, 499-561.
- NUNBERG, J.H., MEADE, J.H., COLE, G., LAWYER, F.C., McCABE, P., SCHWEICKART, V., TAL, R., WITTMAN, V.P., FLATGAARD, J.E. and INNIS, M.A. (1984): Molecular cloning and characterisation of a glucoamylase gene of *Aspergillus awamori*.

- Mol. Cell. Biol. 4, 2306-2315.
- ORR-WEAVER, T.L., SZOSTAK, J.W. and ROTHSTEIN, S.J. (1981): Yeast transformation: A model system for the study of recombination. Proc. Natl. Acad. Sci. USA 78, 6354-6358.
- 5 OTENG-GYANG, K., MOULIN, G., GALZY, P. (1981): Zeitschrift allg. Mikrobiol. 21, 537-544.
- RIGBY, P.W.J., DIEKMANN, M., RHODES, C. and BERG, P. (1977): Labelling desoxyribonucleic acid to high specific activity in vitro by nick translation with DNA-polymerase I. J. Mol. Biol. 113, 237-251.
- 10 ROTHSTEIN, S.J., LAZARUS, C.M., SMITH, W.E., BAULCOMBE, D.C. and GATENBY, A.A. (1984): Secretion of a wheat alpha-amylase expressed in yeast. Nature 308, 662-665.
- RUSSELL, D.W. and HALL, B.D., (1983) J. Biol. Chem. 258, 143-149.
- RUSSELL, D.W., SMITH, M., WILLIAMSON, V.M. and YOUNG, E.T. (1983)
- 15 J. Biol. Chem. 258, 2674-2682.
- SCHATZ, G. (1979): Meth. Enzymol. 1 VI, 40-50.
- SHERMAN, F., FINK, G.R. and HICKS, J.B. (1983): Methods in yeast genetics. In C.S.H. Laboratory, C.S.H., New York, 119.
- 20 SILLS, A.M. and STEWART, G.G. (1982): Production of amylolytic enzymes by several yeast species. J. Inst. Brew. 88, 313-316.
- SILLS, A.M., PANCHAL, C.J., RUSSELL, I. and STEWART, G.G. (1983 a): Genetic manipulation of amylolytic enzyme production of yeast. Gene Expression in Yeast. Proceedings of the Alko Yeast Symposium Helsinki 1983, ed. by M. Korhola and E. Väisänen.
- 25 Foundation for Biotechnical and Industrial Fermentation Research 1, 209-228.
- SILLS, A.M., RUSSELL, I. and STEWART, G.G. (1983 b): The production of yeast amylases in the brewing low carbohydrate beer. EBC Congress, 377-384.
- SILLS, A.M., SAUDER, M.E. and STEWART, G.G. (1984 a): Isolation and characterisation of the amylolytic
- 30 system of Schwanniomyces castellii. J. Inst. Brew. 90, 311-314.
- SILLS, A.M., ZYGORA, P.S.J. and STEWART, G.G. (1984 b): Characterisation of Schwanniomyces castellii mutants with increased productivity of amylases. Appl. Microbiol. Biotechnol. 20, 124-128.
- 35 SIMOES-MENDES, B. (1984): Purification and characterisation of the extracellular amylases of the yeast Schwanniomyces alluvius. Can. J. Microbiol. 30, 1163-1170.
- SOUTHERN, E.M. (1975): Detection of specific sequences among DNA fragments separated by gel electrophoresis. J. Mol. Biol. 98, 503-517.
- 40 STRASSER, A.W.M. (1982): Biochemische und biophysikalische Studien an der in Hefe überproduzierten Hefe Tryptophan-synthetase. Dissertation, Universität Basel.
- STRUHL, K., STINCHCOMB, D.T., SCHERER, S. and DAVIES R.W. (1979): High-frequency transformation of yeast: Autonomous replication of hybrid DNA molecules.
- 45 Proc. Natl. Acad. Sci. USA 76, 1035-1039.
- THOMSEN, K.K. (1983): MOUSE α -amylase synthesised by Saccharomyces cerevisiae is released into the culture medium. Carlsberg Res. Commun. 48, 5, 545-555.
- TOUCHSTONE, J.C. and DOBBINS, M.F., (1978): Practice of thin layer chromatography. A Wiley-Interscience publication. John Wiley & Sons.
- 50 TSAI, H., TSAI, J.H.J. and YU, P.H. (1973): Effects of yeast proteinase and its inhibitors on the inactivation of tryptophane synthetase from Saccharomyces cerevisiae and Neurospora crassa. Eur. J. Biochem. 40, 225-232.
- WALZ, A., RATZKIN, B. and CARBON, J. (1978): Control of expression cloned yeast (Saccharomyces cerevisiae) gene TRP5 by a bacterial insertion element (IS2).
- 55 Proc. Natl. Acad. Sci. USA 75, 6172-6176.
- WILSON, J.J. and INGLEDEW, W.M. (1982): Isolation and characterisation of Schwanniomyces alluvius amylolytic enzymes.

- Appl. Environ. Microbiol. 44, 301-307.
 YAMASHITA, I. and FUKUI, S. (1983): Molecular cloning of glucoamylase-producing gene in the yeast *Saccharomyces*.
 Agric. Biol. Chem. 47, 2689-2692.
 5 ZALKIN, H. and YANOFSKY, C. (1982): Yeast gene TRP5: Structure, function, regulation.
 J. Biol. Chem. 257, 1491-1500.

Claims

- 10 1. A method for producing polypeptides, preferably amylolytic enzymes, by culturing a host microorganism under suitable conditions and recovering of said polypeptides in a manner known per se, characterised in that the host microorganism, as a result of recombinant DNA technology, has received DNA sequences from a donor yeast comprising the coding sequences for said polypeptides and is capable of expressing said polypeptides.
 15 2. A method according to claim 1, characterised in that the said polypeptides are α -amylase and/or glucoamylase.
 3. A method according to one of the claims 1 or 2, characterised in that the host microorganism is selected from a group consisting of yeasts of the genera *Saccharomyces*, *Schizosaccharomyces*, *Schwanniomyces* or *Kluyveromyces*, or bacteria of the genus *Zymomonas* or moulds of the genus *Aspergillus*.
 20 4. A method according to claim 3, characterised in that the yeast is *Saccharomyces cerevisiae*.
 5. A method according to one of the claims 1 to 4 characterised in that the donor yeast is selected from a group consisting of the genera *Saccharomycopsis*, *Lipomyces*, *Pichia*, *Schwanniomyces*, *Debaryomyces* or *Saccharomyces*.
 25 6. A method according to claim 5 characterised in that the donor yeast is *Schwanniomyces castellii* ATCC26076.
 7. A method according to one of the claims 1 to 6, characterised in that the transfer of the DNA sequences from the donor yeast into the host microorganism is carried out by transformation.
 8. A method according to claim 7, characterised in that the transformation is carried out by a vector, preferably a DNA fragment, a plasmid or a cosmid.
 30 9. A method according to claim 8, characterised in that the vector contains homologous DNA, preferably derived from *Schwanniomyces*, leading to integration into the host genome.
 10. A method according to one of the claims 8 or 9 characterised in that the vector contains one or more DNA sequences which control the function of replication and maintenance within the *Schwanniomyces* cells, preferably the autonomously replicating DNA sequences ARS, most preferably the DNA sequences SwARS originating from *Schwanniomyces*.
 35 11. A method according to one of the claims 8 to 10 characterised in that the vector is pCJD5-AMG1 and/or YRpJD2- α .
 12. A method according to one of the claims 7 to 11, characterised in that the host yeast cells before the transformation procedure are subjected to fast freezing, which preferably is carried out with dry ice acetone at temperatures of about -70° to -80°C.
 40 13. A method according to one of the claims 7 to 12 characterised in that the transformation is carried out during thawing and rapid agitation of the frozen yeast cells.
 14. A method according to one of the foregoing claims characterised in that the host yeast is specifically glycosylating the secreted polypeptide.
 45 15. A method according to one of the foregoing claims characterised in that the production of α -amylase is used as marker in *Kluyveromyces lactis*, *Saccharomyces cerevisiae* and *Schizosaccharomyces pombe* for transformation under non-selective conditions.
 16. An α -amylase produced according to the method of one of the claims 1 to 15.
 50 17. An α -amylase according to claim 16, characterised in that it has its temperature optimum at about 37°C and is inactivated at about 60°C.
 18. An α -amylase according to claims 16 or 17, characterised in that it is encoded by the DNA sequence depicted in Figure 2 and all modifications of said DNA sequence which retain the function of the encoded α -amylase.
 55 19. A glucoamylase produced according to the method of one of the claims 1 to 15.
 20. A glucoamylase according to claim 19 characterised in that it has its temperature optimum at about 50°C, and is inactivated at about 60°C and is capable of hydrolysing barley starch completely into glucose.

21. A microorganism, preferably useful as a host microorganism in a method according to one of the claims 1 to 15, characterised in that it is capable of expressing an α -amylase according to claims 16 to 18 and/or a glucoamylase according to claims 19 or 20 as a result of having received DNA sequences from a donor yeast comprising the coding sequences for said α -amylase and/or glucoamylase by recombinant DNA technology.
22. A microorganism according to claim 21, characterised in that it is selected from a group consisting of yeasts of the genera Saccharomyces, Schizosaccharomyces, Schwanniomyces, or Kluyveromyces or bacteria of the genus Zymomonas or moulds of the genus Aspergillus.
23. A yeast cell according to claim 22, characterised in that it is Saccharomyces cerevisiae.
24. A yeast cell according to claim 22, characterised in that it is of the genus Schwanniomyces, preferably the species Schwanniomyces alluvius.
25. A yeast cell according to claim 23 characterised in that it is useful in brewing and baking processes, the food industry, and fermentation processes.
26. A yeast cell according to claim 23 characterised in that it is useful for complete fermentation of starch into ethanol.
27. A yeast cell according to claim 23 characterised in that it is preferably useful in producing special beer by being capable of fermenting dextrins.
28. A vector carrying foreign DNA sequences which are coding for a polypeptide, characterised in that said polypeptides are amylolytic enzymes, preferably α -amylase and/or glucoamylase, wherein the vector is preferably useful in a process of one of the claims 1 to 15.
29. A vector according to claim 28, characterised in that it can be used as a shuttle vector for Escherichia coli -Saccharomyces cerevisiae -Schwanniomyces alluvius -Schizosaccharomyces pombe.
30. A vector according to claims 28 or 29, characterised in that the vector is pCJD5-AMG1 or YRpJD2- α .
31. A vector according to claims 28 or 29, characterised in that it is selected from a group consisting of the plasmids YRpJD2, pCJD5-1 or pCJD5-2.
32. A DNA sequence characterised in that it is coding for a polypeptide, preferably an amylolytic enzyme, most preferably an α -amylase and/or glucoamylase, and is capable of expressing said polypeptide in a suitable host microorganism, preferably a microorganism according to one of the claims 21 to 27 and is preferably useful in a method according to one of the claims 1 to 15.
33. A DNA sequence according to claim 32 characterised in that it has a sequence as depicted in Figure 2.
34. A DNA sequence according to claim 32 characterised in that it has a sequence as depicted in Figure 17.
35. A DNA sequence according to claims 32 or 33, characterised in that it is coding for a α -amylase according to one of the claims 16 to 18, and is obtained by recombinant DNA technology from natural and/or cDNA and/or chemically synthesised DNA.
36. A DNA sequence according to claim 35, characterised in that it comprises the DNA sequence as depicted in Figure 2, encoding the α -amylase, the amino acid sequence of which is depicted in Figure 2.
37. A DNA sequence according to claims 32 or 34 characterised in that it is coding for a glucoamylase according to one of the claims 19 or 20 and is obtained by recombinant DNA technology from natural and/or cDNA or chemically synthesised DNA.
38. A DNA sequence according to claim 37 characterised in that it comprises the DNA sequence as depicted in Figure 17, encoding the glucoamylase, the amino-acid sequence of which is depicted in Figure 17.
39. A combination of DNA sequences characterised in that it comprises a structural gene encoding an amylolytic enzyme and one or more DNA sequences capable of regulation of the expression of the structural gene in a microorganism, preferably, a microorganism according to one of the claims 21 to 27.
40. A combination of DNA sequences according to claim 39, characterised in that it comprises a structural gene encoding an amylolytic enzyme of a yeast.
41. A combination of DNA sequences according to claim 40, characterised in that the structural gene is encoding amylolytic enzymes of the genus Schwanniomyces, preferably of the species Schwanniomyces castellii.
42. A combination of DNA sequences according to one of the claims 39 to 41, characterised in that said structural gene, coding for an amylolytic enzyme is encoding an α -amylase.
43. A combination of DNA sequences according to claim 42 characterised in that the structural gene has the DNA sequence as depicted in Figure 2, and is encoding the polypeptide having the amino acid sequence which is depicted in Figure 2.

44. A combination of DNA sequences characterised in that the structural gene has the DNA sequence as depicted in Figure 17 and is encoding a polypeptide having the amino acid sequence as also depicted in Figure 17.
45. A combination of DNA sequences according to one of the claims 39 to 44 characterised in that the DNA sequence comprises all modifications of the DNA sequences retaining their coding function for an amyolytic enzyme.
46. A combination of DNA sequences according to one of the claims 39 to 45 characterised in that the structural gene contains DNA sequences derived from the structural gene coding for α -amylase or glucoamylase which modify said protein product while retaining its functions in such a way that said product is secreted into the cell wall or the culture medium.
47. A combination of DNA sequences according to claim 46 characterised in that it is contained in a host microorganism according to one of the claims 21 to 27, said host microorganism being suitable in a process for preparing a polypeptide according to one of the claims 1 to 15.
48. A beer manufacturing process characterised in that a yeast is used which has received by recombinant DNA technology DNA sequences from a donor yeast which encodes amyolytic enzymes, preferably α -amylase and/or glucoamylase, wherein the yeast used is preferably Saccharomyces cerevisiae.
49. A beer manufacturing process according to claim 48 characterised in that a yeast being capable of expressing an α -amylase and/or a yeast being capable of expressing glucoamylase or a yeast being capable of expressing α -amylase and glucoamylase is used.
50. A process according to claims 48 or 49 characterised in that the manufacturing process is adapted for the production of a special beer, preferably low carbohydrate beer, low calorie beer or diet beer.
51. A process according to one of the claims 48 to 50 characterised in that the manufacturing process is adapted for the production of beers containing levels of aroma components comparable to beers obtained by fermentation of high glucose worts.
52. A process according to one of the claims 48 to 51 characterised in that the process is decreased in time due to a more rapid decrease in apparent extract (Extrakt scheinbar = Es) at the end of the fermentation process.
53. A process according to one of the claims 48 to 52 characterised in that the manufacturing process is adapted for the production of a beer from a high gravity wort, the carbohydrate level of which can be regulated by a predetermined fermentation time and dilution thereafter.
54. A process according to one of the claims 48 to 53 characterised in that the manufacturing process is adapted for the fermentation of substrates, the constituents of which are for a considerable part suspended hydrated carbohydrates.
55. A process according to one of the claims 48 to 54 characterised in that the genetically engineered yeast is brought into contact with the fermented wort subsequently to main fermentation, but prior to conditioning.
56. An ethanol manufacturing process characterised in that a yeast is used which has received by recombinant DNA technology DNA sequences from a donor yeast which encodes amyolytic enzymes, preferably α -amylase and/or glucoamylase, wherein the yeast used is preferably Saccharomyces cerevisiae.
57. A fermentation process characterised in that a yeast is used which has received by recombinant DNA technology DNA sequences from a donor yeast which encodes amyolytic enzymes, preferably α -amylase and/or glucoamylase, wherein the yeast used is preferably Saccharomyces cerevisiae according to claim 23 and wherein said fermentation process is adapted for the production of beverages produced from fermented or distilled fermented substrates containing suspended hydrated starch.
58. A process for the production of baker's yeast characterised in that a yeast is used which has received by recombinant DNA technology DNA sequences from a donor yeast which encodes amyolytic enzymes, preferably α -amylase and/or glucoamylase, wherein the yeast used is preferably Saccharomyces cerevisiae according to claim 23.
59. A process of preparing bread in a manner known per se, characterised in that a yeast according to claim 58 is used.
60. A process for the production of biomass characterised in that a yeast is used which has received by recombinant DNA technology DNA sequences from a donor yeast which encodes amyolytic enzymes, preferably α -amylase and/or glucoamylase, wherein the yeast used is preferably Saccharomyces cerevisiae according to claim 23.

0 260 404

61. A transformation system for the transfer of DNA sequences from a donor microorganism to a host microorganism, characterised in that the host microorganism is subjected to fast freezing before adding the DNA, preferably that the fast freezing is carried out with dry ice acetone at temperatures of about -70°C to -80°C .

5

10

15

20

25

30

35

40

45

50

55

FIG. 1




FIG. 2A

10 20 30 40 50 60
 GGTACCTGAGCTAAATTTAGAACCGGCTATAGATCCGCTTGTCTAAAGAAGAGATAATGA
 70 80 90 100 110 120
 AGAAAACAATTAACCGAGCACTCTTATTAAGTTTTTCTATTTTCTTTTGCTCCTACTT
 130 140 150 160 170 180
 CAATAATTTATCTAAATTGTATTGTGCGTTAGATCAGAATGTACTGATAACAGAGAGAT
 190 200 210 220 230 240
 TATCATACACTTGTGGATTTCAAAGGCGGAATCAAAGCATACGTAGTCAAACCCTTGG
 250 260 270 280 290 300
 TTATTTGATGCAATTAAGGTTGTAGTCGTTCTTACCGATCCATCATTATACCCACACGG
 310 320 330 340 350 360
 TTTCAATGATGATGAGGTGTTTCAATAGTGAAGTACAATGAATGTTTTGGTAATGCTGTAT
 370 380 390 400 410 420
 GTGGATCAGTAATTATGTTAAACAATTAAGTCTGAAAATTTATTAATAATTTACCTACAA
 430 440 450 460 470 480
 ATTAAGCCGAAATCCAATCGAAGGTGCCGCCAGCTGGTGTATAAATTACTCTTGAAATT
 490 500 510 520 530 540
 CAAGTTGAACGTTGATCTCTCTAAAGCAAAGCTGTTATTCTACAATACTAAATAAAATAA
 550 560 570 580 590 600
 AAGCAAGACATGAGATTTTCAACTGAAGGATTTACAAGTAAAGTTGTTGCAGCAATTTTA
 MetArgPheSerThrGluGlyPheThrSerLysValValAlaAlaIleLeu
 610 620 630 640 650 660
 GCATTCTCAAGATTGGTATCTGCTCAACCGATTATTTTGGACATGAGAGATGTTAGCTCG
 AlaPheSerArgLeuValSerAlaGlnProIleIlePheAspMetArgAspValSerSer
 670 680 690 700 710 720
 TCAGCTGATAAATGGAAAGACCAATCGATTTATCAAATCGTTACTGATAGGTTTGCCAGA
 SerAlaAspLysTrpLysAspGlnSerIleTyrGlnIleValThrAspArgPheAlaArg
 730 740 750 760 770 780
 TCTGATGGCTCGACCACAGCTGACTGTTTAGTGAGTGATCGCAAGTACTGTGGTGGATCT
 SerAspGlySerThrThrAlaAspCysLeuValSerAspArgLysTyrCysGlyGlySer
 790 800 810 820 830 840
 TATAAAGGGATTATCGACAAGTTGGATTATATTCAAGGTATGGGTTTCACTGCGATCTGG
 TyrLysGlyIleIleAspLysLeuAspTyrIleGlnGlyMetGlyPheThrAlaIleTrp
 850 860 870 880 890 900
 ATCTCCCCAGTTGTTGAGCAAATTCCTGACAATACTGCTTATGGTTATGCTTACCATGGT
 IleSerProValValGluGlnIleProAspAsnThrAlaTyrGlyTyrAlaTyrHisGly

FIG. 2 B

910 920 930 940 950 960
 TATTGGATGAAAAATATTGATGAATTGAACACTAATTTTGGTACCGCTGATGAATTGAAA
 TyrTrpMetLysAsnIleAspGluLeuAsnThrAsnPheGlyThrAlaAspGluLeuLys

970 980 990 1000 1010 1020
 CAATTAGCTAGCGAATTGCATTTTCAGAAGCATGTATGATGGTCGACGATGTTACAACCAT
 GlnLeuAlaSerGluLeuHisPheArgSerMetTyrAspGlyArgArgCysTyrAsnHis

1030 1040 1050 1060 1070 1080
 TATGCTTGGAAACGGAGATGGTTCAAGCGTAGATTATTCTAGTTTCACTCCATTCAATCAA
 TyrAlaTrpAsnGlyAspGlySerSerValAspTyrSerSerPheThrProPheAsnGln

1090 1100 1110 1120 1130 1140
 CAATCTTACTTCCACGATTATTGTTTGATTACAAATTATAATGATCAAACCAATGTTGAA
 GlnSerTyrPheHisAspTyrCysLeuIleThrAsnTyrAsnAspGlnThrAsnValGlu

1150 1160 1170 1180 1190 1200
 GATTGTTGGGAABGTGATACTGAAGTCTCCCTTCCAGATTTAAGTACCGAGGATAATGAA
 AspCysTrpGluGlyAspThrGluValSerLeuProAspLeuSerThrGluAspAsnGlu

1210 1220 1230 1240 1250 1260
 GTTATAGGAGTATTTCAAACCTTGGGTGTCAGATTTTGTCAAACTATTCAATCGATGGT
 ValIleGlyValPheGlnThrTrpValSerAspPheValGlnAsnTyrSerIleAspGly

1270 1280 1290 1300 1310 1320
 TTAAGAATTGATAGTGCAAAGCACGTAGATACCGCTTCATTACGAAGTTTGAGGACGCT
 LeuArgIleAspSerAlaLysHisValAspThrAlaSerLeuThrLysPheGluAspAla

1330 1340 1350 1360 1370 1380
 TCTGGTGTTTATAACTTAGGTGAAGTTTATCAAGGAGATCCAACCTTATACTTGTCCATAT
 SerGlyValTyrAsnLeuGlyGluValTyrGlnGlyAspProThrTyrThrCysProTyr

1390 1400 1410 1420 1430 1440
 CAGAATTATATGAAAGGAGTTACCAACTATCCATTATACTATCCAGTATATAGATTCTTC
 GlnAsnTyrMetLysGlyValThrAsnTyrProLeuTyrTyrProValTyrArgPhePhe

1450 1460 1470 1480 1490 1500
 AGTGATACTTCGGCGACTTCCAGTGAGTTAAGTTCAATGATCTCCACGTTACAGTCATCT
 SerAspThrSerAlaThrSerSerGluLeuThrSerMetIleSerThrLeuGlnSerSer

1510 1520 1530 1540 1550 1560
 TGTTCCGGACGTCTCTTTGTTGGGAACTTTATTGAAAACCATGACCAAGTTAGATTTCCT
 CysSerAspValSerLeuLeuGlyAsnPheIleGluAsnHisAspGlnValArgPhePro

1570 1580 1590 1600 1610 1620
 TCAGTTACCTCAGACACATCCTTGATTAGAATGACATGGCTTTTATAATTTTGGGTGAT
 SerValThrSerAspThrSerLeuIleLysAsnAspMetAlaPheIleIleLeuGlyAsp

1630 1640 1650 1660 1670 1680
 GGTATCCCAATTATTTATTATGGCCAAGAACAAGGTCTCAATGGTGGTTCCGATCCTGCC
 GlyIleProIleIleTyrTyrGlyGlnGluGlnGlyLeuAsnGlyGlySerAspProAla

1690 1700 1710 1720 1730 1740
 AATAGAGAAGCTTTATGGTTAAGTGGATATAATACCGATTGAGAATACTACGAGCTAATC
 AsnArgGluAlaLeuTrpLeuSerGlyTyrAsnThrAspSerGluTyrTyrGluLeuIle

1750 1760 1770 1780 1790 1800
 AGTAACTAAATCAAATAAGAAATCAAGCTATTAAGAAGGATTCTGCCTATTCAACTTAC
 SerLysLeuAsnGlnIleArgAsnGlnAlaIleLysLysAspSerAlaTyrSerThrTyr

FIG. 2C

1810 1820 1830 1840 1850 1860
 AAATCCTCAGTTGTTTCTTCTTCAGACCACTATATAGCCACTAGGAAGGGTAGCGATGCT
 LysSerSerValValSerSerSerAspHisTyrIleAlaThrArgLysGlySerAspAla

1870 1880 1890 1900 1910 1920
 AATCAATTGATTTCATTTTAAATAATTTAGGTTCAAACGGGTCACAGGATATTACTGTC
 AsnGlnLeuIleSerIlePheAsnAsnLeuGlySerAsnGlySerGlnAspIleThrVal

1930 1940 1950 1960 1970 1980
 AGCAACACCGGCTATTCTAGTGGTGATAAAGTTATCGATATTATTTCTTGCAATTCCGTT
 SerAsnThrGlyTyrSerSerGlyAspLysValIleAspIleIleSerCysAsnSerVal

1990 2000 2010 2020 2030 2040
 TTAGCTGGTGACTCCGGAAGCTTATCTGTATCAATTTCTGGTGGAATGCCACAAGTTTAC
 LeuAlaGlyAspSerGlySerLeuSerValSerIleSerGlyGlyMetProGlnValTyr

2050 2060 2070 2080 2090 2100
 GCTCCGTCCTCTGTTCTTTGCGGATCTGGCATCTGCAATCAATAGATTGATCCAGCGCTA
 AlaProSerSerValLeuSerGlySerGlyIleCysAsnGlnEnd

2110 2120 2130 2140 2150 2160
 ACCCTTTTTTTAGCAACGACAAGTTTATTTAGAAAAAGTTTTCTAAGAATGGTCAAAAC

2170 2180 2190 2200 2210 2220
 AAGTTCTTATTACTTCTATGTCCCTGGATATCTGTTTTCAATGTTCTCTGACTCCACATT

2230 2240 2250 2260 2270 2280
 CCTCATGTTTAGTTCTCTATTTTTTGTGATCTTCTAAGTTTTTTTATTCTTAATTTTAA

2290
 TCCAAAAGTT

1

5

FIGURE 3

10 PEPTIDE 1 : gly-gly-asn-val-leu-pro-thr-gln-glu-pro-gly-tyr-thr-
val-ala-glu-ser-arg

PEPTIDE 2 : ala-thr-ser-tyr-pro-val-gly-phe-asp-val-ser-cys-ala-
ser-glu-trp-lys

15

PEPTIDE 3 : gly-val-phe-pro-gly-ala-gly-lys-glu-glu-val-tyr-tyr-
asp-trp-tyr-thr-gln-arg

PEPTIDE 4 : trp-gly-tyr-asp-thr-ile-glu-lys

20


PEPTIDE 5 : thr-leu-phe-ala-asn-asp-val-gly-asp-pro-ile-asp-gly-
asn-ile-tyr-gly-val-his-pro-val-tyr-leu-asp-gln-arg

25

30


35

FIG. 4 A




- cell density (log OD 600)
- total α -amylase activity
- ▲—▲ specific activity (activity/OD 600)



FIG. 4 B



- cell density (log OD 600)
- total α -amylase activity
- ▲—▲ specific activity (activity/OD 600)

FIG. 4 C



  cell density (log OD 600)




  total α -amylase activity


FIG. 4 D



- cell density (log OD 600)
- total α -amylase activity
- ▲—▲ specific activity (activity/OD 600)

FIG. 4 E

glucose concentration



- NGA23 - YRpJD2 , 2 % Maltose
- NGA23 - YRpJD2-α , 2 % Maltose
- ▲-▲ NGA23 - YRpJD2 , 2 % Glucose
- ▼-▼ NGA23 - YRpJD2-α , 2 % Glucose






FIG. 5



Fig.6



1707-81

FIG. 8 (2)







B: Bam HI Bg: Bgl II C: Cla I S: Sal I

120000

FIG. 7



A FIG. 6 (1)









FIG. 8 (5)





B**FIG. 8 (6)**

E: EcoRI

B: BamHI

pBR322

S. cerevisiae

C

B: Bam HI Bg: Bgl II C: Cla I S: Sal I
X: Xba I

pBR322

S. castellii

K. lactis

S. cerevisiae

27 07 87


Fig.9⁽¹⁾

Fig.9 (2)



C

D Hybridizingprobe : YRpJD1






Fig.10 (1)



15h

Fig.10 (2)






Fig.11 (1)









Fig.12

Fig.13 (1)

A



— pBR322

— λ

□ *S. castellii*

■ *S. cerevisiae*

E: *EcoRI*

P: *PvuII*

S: *SalI*

B: *BamHI*

Bg: *BglII*

C: *ClaI*




Fig.13 (2)

NOV 87

FIG. 14





FIG. 15 (1)



A : *Ava* I
 Bg : *Bgl* II
 B : *Bam* HI
 E : *Eco* RI
 N : *Nru* I
 P : *Pvu* II
 Ps : *Pst* I
 S : *Sal* I
 X : *Xho* I

————— pBR 322
 ■■■■■ *Saccharomyces cerevisiae*
 ▨▨▨▨ *Kluyveromyces fragilis*
 □□□□ *Schwanniomyces castellii*
 ▨▨▨▨ 2 μ DNA
 ————— λ DNA




FIG. 15 (2)





FIG. 15 (3)





FIG. 16





Fig. 16. Inactivation of α -amylase at different temperatures

FIGURE 17A

-310

AGATCTACATTTTAAACCCAG

-250

TCTACTCCAGATATTGGAGTATAACCCATTCTTACCGTTATATCCATGACCCGCATCGA

-200

AATTTTCAAAGGATTTTCGAGGAAATTCTTTCCTAAATACGAAGTGTTATTGGTGATTCA

-150

ATTACTACGSAACTACTCATATGGTAGTAGAGTTGGTGAATGTAGCGCAATTGTAATT

-100

GCGAAGTTATAGTAATAGTTTGGCAAACCTGGAGAATTTTCATTATTGGGAAAATATAAA

-50

-1

TAAAGGCAAGTATCCATTGAAATTTTAAATGAACTCATGACTGTATTATAACAAGCAAG

FIGURE 17B

10 20 30 40 50 60
 ATGATTTTTCTGAAGCTGATTAAAAGTATAGTAATTGGTTTGGGATTAGTTAGTGCTATC
 MetIlePheLeuLysLeuIleLysSerIleValIleGlyLeuGlyLeuValSerAlaIle

70 80 90 100 110 120
 CAAGCAGCCCCCTGCCTCTTCGATTGGATCTAGTGCTTCAGCATCTAGTTCAAGTGAGAGT
 GlnAlaAlaProAlaSerSerIleGlySerSerAlaSerAlaSerSerSerGluSer

130 140 150 160 170 180
 TCTCAGGCTACAATCCCAATGATGTAACATTAGGTGTTAAACAAATTCCTAATATCTTT
 SerGlnAlaThrIleProAsnAspValThrLeuGlyValLysGlnIleProAsnIlePhe

190 200 210 220 230 240
 AATGACTCTGCTGTCGATGCTAATGCAGCTGCTAAAGGGTATGACTTGGTAAATGTTACT
 AsnAspSerAlaValAspAlaAsnAlaAlaAlaLysGlyTyrAspLeuValAsnValThr

250 260 270 280 290 300
 AATACTCCAAGAGGATTAACCGGTATCTTAAATTAAGAAGCTACCAATATTTATGGT
 AsnThrProArgGlyLeuThrGlyIleLeuLysLeuLysGluAlaThrAsnIleTyrGly

310 320 330 340 350 360
 TATGATTTTGATTATTTAACTTAACTGTTGAATACCAAGCTGATACCAGATTAAACGTT
 TyrAspPheAspTyrLeuAsnLeuThrValGluTyrGlnAlaAspThrArgLeuAsnVal

370 380 390 400 410 420
 CATATTGAACCAACTGATTTATCTGATGTATTTGTTTTACCAGAGCATTAGTTGTTAA
 HisIleGluProThrAspLeuSerAspValPheValLeuProGluHisLeuValValLys

430 440 450 460 470 480
 CCACTGGTGGAAGGTGATGCACAATCTTATAACTTCGACAATTCGGATTGGTTTTCGAA
 ProLeuValGluGlyAspAlaGlnSerTyrAsnPheAspAsnSerAspLeuValPheGlu

490 500 510 520 530 540
 TACTCTAATACTGACTTCTCCTTTGAAGTTATTAGATCATCTACTAAAGAAGTTTTATT
 TyrSerAsnThrAspPheSerPheGluValIleArgSerSerThrLysGluValLeuPhe

550 560 570 580 590 600
 TCTACTAAAGGTAATCCATTGGTTTTTCAAATCAATTCATTCAATTCATTCGTCATTG
 SerThrLysGlyAsnProLeuValPheSerAsnGlnPheIleGlnPheAsnSerSerLeu

610 620 630 640 650 660
 CCAAAGAACCATGTTATTACTGGTCTTGGTGAATCTATTACGGTTTTAGTTAACGAACCA
 ProLysAsnHisValIleThrGlyLeuGlyGluSerIleHisGlyLeuValAsnGluPro

670 680 690 700 710 720
 GGTAGCGTTAAACATTATTTGCTAATGATGTTGGTGATCCAATCGATGGTAATATTTAT
 GlySerValLysThrLeuPheAlaAsnAspValGlyAspProIleAspGlyAsnIleTyr

730 740 750 760 770 780
 GGTGTCCATCCAGTTTATCTTGATCAAGATATGACACTGAAACTACCCATGCTGTTTAT
 GlyValHisProValTyrLeuAspGlnArgTyrAspThrGluThrThrHisAlaValTyr

790 800 810 820 830 840
 TGGAGAACTTCTGCTATTCAAGAAGTATTAATCGGTGAGGAATCTATTACTTGGAGAGCT
 TrpArgThrSerAlaIleGlnGluValLeuIleGlyGluGluSerIleThrTrpArgAla

850 860 870 880 890 900
 CTTTCAGGTGTTATTGATTTATACTTCTTTAGTGGTCTACACCAAAGATGCCATTCAA
 LeuSerGlyValIleAspLeuTyrPhePheSerGlyProThrProLysAspAlaIleGln

FIGURE 17C

910 920 930 940 950 960
 CAGTATGTCAAAGAGATTGGTTTACCAGCTTTCCAACCATACTGGTCGTTAGGTTACCAT
 GlnTyrValLysGluIleGlyLeuProAlaPheGlnProTyrTrpSerLeuGlyTyrHis

970 980 990 1000 1010 1020
 CAATGTAGATGGGGTTACGATACTATCGAAAAATTATCTGAAGTTGTTGAAAACCTTCAAG
 GlnCysArgTrpGlyTyrAspThrIleGluLysLeuSerGluValValGluAsnPheLys

1030 1040 1050 1060 1070 1080
 AAATTTAATATTCCATTAGAACTATCTGGTCAGACATTGATTACATGGACTCTTATAAA
 LysPheAsnIleProLeuGluThrIleTrpSerAspIleAspTyrMetAspSerTyrLys

1090 1100 1110 1120 1130 1140
 GATTTCACTTATGATCCACACAGATTCCTACTAGATGAATATCGTAAATTCCTTGATGAG
 AspPheThrTyrAspProHisArgPheProLeuAspGluTyrArgLysPheLeuAspGlu

1150 1160 1170 1180 1190 1200
 TTGCACAAAAATAATCAACACTATGTTCTATTTTGGATGCTGCTATTTACGTTCCAAAC
 LeuHisLysAsnAsnGlnHisTyrValProIleLeuAspAlaAlaIleTyrValProAsn

1210 1220 1230 1240 1250 1260
 CCAACAATGCTACGGATAACGAATACCAACCTTTCCACTATGGTAATGAAACCGATGTC
 ProAsnAsnAlaThrAspAsnGluTyrGlnProPheHisTyrGlyAsnGluThrAspVal

1270 1280 1290 1300 1310 1320
 TTCTTAAAGAATCCAGATGGTTCATTATATATTGGTGTGTTTGGCAGGTTACACTGTTT
 PheLeuLysAsnProAspGlySerLeuTyrIleGlyAlaValTrpGlnValThrLeuPhe

1330 1340 1350 1360 1370 1380
 TCCAGATTTCTTAGCAGAAACATTTCAGATATGGATAAAGTCATTAAAGATTGGTATGAA
 SerArgPheLeuSerArgLysHisSerAspMetAspLysValIleLysAspTrpTyrGlu

1390 1400 1410 1420 1430 1440
 TTAACCTCTTTTGATGGTATTTGGGCTGATATGAATGAAGTCTCATCTTCTGTGTTGGT
 LeuThrProPheAspGlyIleTrpAlaAspMetAsnGluValSerSerPheCysValGly

1450 1460 1470 1480 1490 1500
 TCTTGTGGTACTGGTAAATACTTCGAAAACCCAGCATATCCTCCATTTACTGTTGGAAGT
 SerCysGlyThrGlyLysTyrPheGluAsnProAlaTyrProProPheThrValGlySer

1510 1520 1530 1540 1550 1560
 AAAGCTACCTCTTATCCAGTTGGTTTCGATGTTTCTAACGCATCTGAATGGAAATCTATT
 LysAlaThrSerTyrProValGlyPheAspValSerAsnAlaSerGluTrpLysSerIle

1570 1580 1590 1600 1610 1620
 CAAAGCTCAATTTCTGCTACTGCTAAGACTTCTTCAACTTCTTCCGTATCGTCGTCTTCA
 GlnSerSerIleSerAlaThrAlaLysThrSerSerThrSerSerValSerSerSerSer

1630 1640 1650 1660 1670 1680
 TCCACAATCGATTATATGAACACTTTAGCTCCAGGTAAAGGTAATATTAATTATCCACCA
 SerThrIleAspTyrMetAsnThrLeuAlaProGlyLysGlyAsnIleAsnTyrProPro

1690 1700 1710 1720 1730 1740
 TATGCTATTTACAACATGCAAGGTGACTCCGATCTTGCTACTCATGCAGTATCTCCAAAT
 TyrAlaIleTyrAsnMetGlnGlyAspSerAspLeuAlaThrHisAlaValSerProAsn

1750 1760 1770 1780 1790 1800
 GCTACACATGCTGATGGTACAGTTGAATATGATATTCACAATCTTTATGGTTACTTGCAA
 AlaThrHisAlaAspGlyThrValGluTyrAspIleHisAsnLeuTyrGlyTyrLeuGln

0 260 404
FIGURE 17D

1810	1820	1830	1840	1850	1860
GAAATGCTACTTATCATGCATTATTGGAAGTTTTTCCTAACAAAGAGACCATTTCATGATT					
GluAsnAlaThrTyrHisAlaLeuLeuGluValPheProAsnLysArgProPheMetIle					
1870	1880	1890	1900	1910	1920
TCCAGATCAACCTTTCCAXGCGCTGGTAAATGGACCGGCCATTGGGGTGGTGACAACACT					
SerArgSerThrPhePro AlaGlyLysTrpThrGlyHisTrpGlyGlyAspAsnThr					
1930	1940	1950	1960	1970	1980
GCTGATTGGGCTTATGCTTACTTCTCTATCCCTCAAGCATTCTCAATGGGTATTGCTGGC					
AlaAspTrpAlaTyrAlaTyrPheSerIleProGlnAlaPheSerMetGlyIleAlaGly					
1990	2000	2010	2020	2030	2040
CTTCCATTCTTTGGTGCCGATGTTTGTGGTTTCAATGGTAATTCTGATTCTGAATTATGT					
LeuProPhePheGlyAlaAspValCysGlyPheAsnGlyAsnSerAspSerGluLeuCys					
2050	2060	2070	2080	2090	2100
TCAAGATGGATGCAATTAGGTTCTTTCTTCCCATTTCTACAGAAACCACAACCTATTTAGGT					
SerArgTrpMetGlnLeuGlySerPhePheProPheTyrArgAsnHisAsnTyrLeuGly					
2110	2120	2130	2140	2150	2160
GCTATTGATCAGGAACCATATGTCTGGGAATCAGTTGCTGAAGCTACTAGAACTTCTATG					
AlaIleAspGlnGluProTyrValTrpGluSerValAlaGluAlaThrArgThrSerMet					
2170	2180	2190	2200	2210	2220
GCCATTAGATACTTATTATTACCATATTACTACACTTTATTACATGAATCTCATACTACT					
AlaIleArgTyrLeuLeuLeuProTyrTyrTyrThrLeuLeuHisGluSerHisThrThr					
2230	2240	2250	2260	2270	2280
GGTTTACCAATCTTAAGAGCTTTCTCGTGGCAATTCCCTAACGATCGTTCCTTAAGTGGT					
GlyLeuProIleLeuArgAlaPheSerTrpGlnPheProAsnAspArgSerLeuSerGly					
2290	2300	2310	2320	2330	2340
GTCGATAACCAATTTTTTGTCTGGTGTATGGTTTAGTTGTTACTCCTGTCTTAGAACCTGGT					
ValAspAsnGlnPhePheValGlyAspGlyLeuValValThrProValLeuGluProGly					
2350	2360	2370	2380	2390	2400
GTTGATAAGGTTAAAGGTGTTTTCCAGGAGCTGGTAAGAGGAAGTTTACTACGACTGG					
ValAspLysValLysGlyValPheProGlyAlaGlyLysGluGluValTyrTyrAspTrp					
2410	2420	2430	2440	2450	2460
TACACCCAAAGAGAAGTTCACCTTTAAAGACGGTAAGAATGAACTTTAGATGCACCATTA					
TyrThrGlnArgGluValHisPheLysAspGlyLysAsnGluThrLeuAspAlaProLeu					
2470	2480	2490	2500	2510	2520
GGTCATATTCCATTACACATTAGAGGTGGTAACGTCTTGCCAACTCAAGAGCCAGGTTAT					
GlyHisIleProLeuHisIleArgGlyGlyAsnValLeuProThrGlnGluProGlyTyr					
2530	2540	2550	2560	2570	2580
ACTGTTGCTGAGTCAAGACAAAATCCATTTGGTTTAATTGTCGCTTTAGATAACGATGGC					
ThrValAlaGluSerArgGlnAsnProPheGlyLeuIleValAlaLeuAspAsnAspGly					
2590	2600	2610	2620	2630	2640
AAAGCTCAAGGTAGCTTATACCTTGATGATGGTGAATCATTAGTAGTAGACTCTTCATTG					
LysAlaGlnGlySerLeuTyrLeuAspAspGlyGluSerLeuValValAspSerSerLeu					
2650	2660	2670	2680	2690	2700
TTGGTTAGTTTCTCTGTTTCTGATAACACATTATCAGCATCTCCATCTGGTGACTATAAA					
LeuValSerPheSerValSerAspAsnThrLeuSerAlaSerProSerGlyAspTyrLys					

FIGURE 17E

2710 2720 2730 2740 2750 2760
GCTGATCAACCTTTAGCTAATGTTACCATCTTAGGGGTTGGCCATAAACCAAAATCAGTT
AlaAspGlnProLeuAlaAsnValThrIleLeuGlyValGlyHisLysProLysSerVal

2770 2780 2790 2800 2810 2820
AAATTTGAAAACGCTAATGTTGATTTACCTACAAGAAATCAACCGTTTTTCGTTACTGGC
LysPheGluAsnAlaAsnValAspPheThrTyrLysLysSerThrValPheValThrGly

2830 2840 2850 2860 2870 2880
TTAGATAAATACACCAAGGATGGTGCATTTTCTAAGGATTTACCATTTACTTGTAATTT
LeuAspLysTyrThrLysAspGlyAlaPheSerLysAspPheThrIleThrTrp

2890 2900 2910 2920 2930 2940
TAACATCCACTTAGTTCAATTCATTCTTTTCTTTTCCCGTGAAATTCTGAATTTGAAA

2950 2960 2970 2980 2990 3000
TTATTTGAATGATATCATTTTAGTTTTCTTCAGCTTATGCTATGTTTATTTTCGATTTTAA

3010 3020 3030 3040 3050 3060
ATGTTAAAAGTTTTTTATGTTTATGTTGTTTTATTGATGTAGTTGATAAAATATAGCAAA

3070 3080 3090 3100 3110 3120
TACATCGAAAAATTTGCGATGAAATTTGCAGCTCATTAGAAATGTAGTCAATCATTAGT

3130 3140 3150 3160 3170 3180
CACATTGGACCACTATATAACAAACAACAATTTCCAAGAAAAATATATGTAAGGATAC

3190 3200 3210 3220 3230 3240
TAGATCATAAATTCTTATTGACTTTGTTTTTTTAAACAATAGTTACATAAGGAATATTCTG

3250 3260 3270 3280 3290 3300
TTTACTACAAAACCATTTGGTCTTGTAAGAAGCAGACGAGGCGTATGTTTGTGGTTGCGG

3310 3320
CCGCAATACTAGTTTACAAAG

THIS PAGE BLANK (USPTO)

12 **EUROPEAN PATENT APPLICATION**

21 Application number: 87110370.1

22 Date of filing: 17.07.87

51 Int. Cl. 4: **C12N 15/00**, **C12P 21/02**,
C12N 9/30, **C12N 9/34**,
C12P 7/06, **C12P 1/00**,
C12C 11/04, **C12N 1/16**,
A21D 8/04, **C12N 1/18**

30 Priority: 21.08.86 EP 86111586

43 Date of publication of application:
23.03.88 Bulletin 88/12

84 Designated Contracting States:
AT BE CH DE ES FR GB GR IT LI LU NL SE

89 Date of deferred publication of the search report:
06.07.88 Bulletin 88/27

71 Applicant: **Heineken Technisch Beheer B.V.**
Burg. Smeetsweg 1
NL-2382 PH Zoeterwoude(NL)


72 Inventor: **Strasser, Alexander**
Zonser Strasse 6
D-4000 Düsseldorf(DE)
 Inventor: **Martens, Feodor Bernard**
Hogenwoerd 13
NL-2311 HE Leiden(NL)
 Inventor: **Dohmen, Jürgen**
Bretonenstrasse 20
D-4005 Meerbusch 1(DE)
 Inventor: **Hollenberg, Cornelius P.**
Chopinstrasse 7
D-4000 Düsseldorf(DE)

74 Representative: **Patentanwälte Grünecker,**
Kinkeldey, Stockmair & Partner
Maximilianstrasse 58
D-8000 München 22(DE)

54 **Amylolytic enzymes producing microorganisms constructed by recombinant DNA technology and their use for fermentation processes.**

57 This invention provides a method for producing amylolytic enzymes by culturing a microorganism, having received as a result of recombinant DNA technology DNA sequences from a donor yeast comprising the coding sequences for the amylolytic enzymes wherein the host microorganism is capable of expressing said amylolytic enzymes. Furthermore, this invention provides microorganisms genetically engineered as to being able to produce and express the amylolytic enzymes, a vector containing the DNA sequences, coding for the amylolytic enzymes and the respective DNA sequences. The said host microorganisms are useful in the production of beer and many fermentation processes, preferably in the production of special

beers.



Xerox Copy Centre



European Patent
Office

EUROPEAN SEARCH REPORT

Application number

EP 87 11 0370

DOCUMENTS CONSIDERED TO BE RELEVANT			
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	CLASSIFICATION OF THE APPLICATION (Int. Cl.4)
X	WO - A - 86 03 778 (BREWING RESEARCH FOUND) * Pages 1,2; page 4, paragraphs 3-6; pages 13,14: section a; page 17, table II,; page 21, section d; page 24, last paragraph, page 26, paragraph 3; claims 4-10 *	1-5, 7,8, 10,14, 19,21- 23,25- 29,32, 37,39, 40,45- 57	C 12 N 15/00 C 12 P 21/02 C 12 N 9/30 C 12 N 9/34 C 12 P 7/06 C 12 P 1/00 C 12 C 11/04 C 12 N 1/16 A 21 D 8/04 C 12 N 1/18
Y		6,9, 11-13, 34, 38,41, 44,58- 60	
			TECHNICAL FIELDS SEARCHED (Int. Cl.4)
X	EP - A - 0 173 668 (SMITHKLINE BECKMAN) * Page 3, lines 9-16; page 5, line 32 - page 6, line 3; page 8, lines 18-23; page 15, lines 9-29; page 19, line 25 - page 22, line 30; claim 30 *	1-4, 7,8, 14,19, 21- 23,28, 29,32, 37,39, 40, 45-57	C 12 N C 12 P C 12 C A 21 D
		-2-	
Place of search THE HAGUE		Date of completion of the search 14-12-1987	Examiner MADDOX
<p>CATEGORY OF CITED DOCUMENTS</p> <p>X : particularly relevant if taken alone Y : particularly relevant if combined with another document of the same category A : technical background O : non-written disclosure P : intermediate document</p> <p>T : theory or principle underlying the invention E : earlier patent document, but published on, or after the filing date D : document cited in the application L : document cited for other reasons & : member of the same patent family, corresponding document</p>			

EPO form 1503 03 82



European Patent
Office

CLAIMS INCURRING FEES

The present European patent application comprised at the time of filing more than ten claims.

- ☐ All claims fees have been paid within the prescribed time limit. The present European search report has been drawn up for all claims.
- ☐ Only part of the claims fees have been paid within the prescribed time limit. The present European search report has been drawn up for the first ten claims and for those claims for which claims fees have been paid,
namely claims:
- ☐ No claims fees have been paid within the prescribed time limit. The present European search report has been drawn up for the first ten claims.

X LACK OF UNITY OF INVENTION

The Search Division considers that the present European patent application does not comply with the requirement of unity of invention and relates to several inventions or groups of inventions,
namely:

1. Claims 1-60: Development of amylolytic yeast by genetic engineering
2. Claim 61: A method for transforming microorganisms

- ☐ All further search fees have been paid within the fixed time limit. The present European search report has been drawn up for all claims.
- ☐ Only part of the further search fees have been paid within the fixed time limit. The present European search report has been drawn up for those parts of the European patent application which relate to the inventions in respect of which search fees have been paid,
namely claims:
- ☒ None of the further search fees has been paid within the fixed time limit. The present European search report has been drawn up for those parts of the European patent application which relate to the invention first mentioned in the claims,
namely claims: 1-60



Eur pean Patent
Office

EUROPEAN SEARCH REPORT

Application number

EP 87 11 0370 -2

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int. Cl.4)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
Y		6,9,11-13,15,34,38,41,44,58-60	
X	-- EP - A - 0 163 491 (BIOTECHNICA) * Page 2, line 24 - page 3, line 26; page 4, lines 10-29; line 30 - page 5, line 12; page 8, lines 10-13; page 10, lines 15-18; page 13, lines 27-32; claims 2,10,12, 15-19 *	1-5,7-10,19,21-23,25-29,32,37,39,40,45-57	
Y		6,9,11-13,15,34,38,41,44,58-60	TECHNICAL FIELDS SEARCHED (Int. Cl.4)
X	-- I.S. PRETORIUS et al.: "Molecular cloning and characterization of the STA2 glucoamylase gene of Saccharomyces diastaticus", & MGG,MOL.GEN. GENET. 1986, 203(1), 29-35 * * Summary *	1-5,7-10,19,21-23,28,29,32,37,39,40,45	
Y		12,13	
X	-- FOOD TECHNOLOGY, vol. 38, no. 2, February 1984, pages 99-106,111; Chicago, US C.J. PANCHAL et al.: "Genetic manipulation of brewing and related yeast strains" * Whole article; in particular page 105, section "Applications" *	1-8,14,19-23,25-29,32,34,37-41,44-57	

EPO Form 1505.3 08.78



European Patent
Office

EUROPEAN SEARCH REPORT

Application number

EP 87 11 0370 -3

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int. Cl.4)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
Y		12,13, 16-18, 33,35, 36,42, 43,58- 60	
O,X	P.G. MEADEN et al.: "A plasmid vector system for the genetic manipulation of brewing strains", & PROC.CONG.-EUR.BREW.CONV. 1985, 20th, 219-26 * Whole document *	1-5,7, 8,19, 21-23, 25-28, 32-37, 39,40, 46-54, 57	
O,Y		6,34, 38,41, 44	TECHNICAL FIELDS SEARCHED (Int. Cl.4)
X	AGRICULTURAL AND BIOLOGICAL CHEMISTRY, vol. 49, no. 10, October 1985, pages 3089-3091, Tokyo, JP; I. YAMASHITA et al.: "Cloning and expression of the Saccharomycopsis fibuligera alpha-amylase gene in Saccharomyces cerevisiae" * Whole article *	1-5,7, 8,14, 16,17, 21-23, 25,26, 28,29, 32,35, 39,40, 42,45- 50,56, 57,60	
X	AGRICULTURAL AND BIOLOGICAL CHEMISTRY, vol. 47, no. 11, November 1983, pages 2689-2692, Tokyo, JP; I. YAMASHITA et al.: "Molecular cloning of a glucoamylase producing gene in the yeast Saccharomyces" * Whole article *	1-5,7,	

EPO Form 1505.3 08.78



DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int. Cl.4)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
X	-- AGRICULTURAL AND BIOLOGICAL CHEMISTRY, vol. 48, no. 7, July 1984, pages 1931-1932, Tokyo, JP; I. YAMASHITA et al.: "Secretion of Saccharomyces diastaticus glucoamylase from Schizosaccharomyces pombe" * Whole article *	8,10-14, 19-23, 28,29, 37,39, 40,45-47	
X	-- NATURE, vol. 308, no. 5960, April 12, 1984, pages 662-65, Reading, Berks, GB; S.J. ROTHSTEIN et al.: "Secretion of a wheat alpha-amylase expressed in yeast" * Page 663, right-hand column *	1-5,7, 8,10, 14,19-23,28, 29,37, 39,40, 45-57 39,42	TECHNICAL FIELDS SEARCHED (Int. Cl.4)
Y	-- GENE, vol. 25, no. 2/3, November 1983, pages 333-341, Elsevier Brussels, BE; R.J. KLEBE et al.: "A general method for polyethylene-glycol induced genetic transformation of bacteria and yeast " * Page 335, section e *	15 12,13	
Y	-- C.J. PANCHAL et al.: "Susceptibility of Saccharomyces spp. and Schwanniomyces spp. to the aminoglycoside antibiotic G418" & APPL.ENVIRON.MICROBIOL. 1984, 47(5), 1164-6 * Left-hand column, paragraph 1 *		
X	-- EP - A - 0 125 615 (LABATT) * Pages 1-4,9,10; page 11, paragraphs 2,3; page 12; page 16,	1-60 16-20	



European Patent
Office

EUROPEAN SEARCH REPORT

Application number

EP 87 11 0370-5-

DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int. Cl. 4)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
Y	last paragraph - page 17, paragraph 1; claims 2-4,8,9,20	12,13, 33-60	
X	-- BIOTECHNOLOGY LETTERS, vol. 5, no. 12, 1983, pages 825-830, M.R. DHAWALE et al.: "Interspecific protoplast fusion of Schwanniomyces yeasts" * Whole article *	24	
X	-- BIOTECHNOLOGY, vol. 4, no. 4, April 1986, pages 311-315, New York, US; S.A. FILHO et al.: "Stable yeast transformants that secrete function- al alpha-amylase encoded by cloned mouse pancreatic cDNA" * Page 314, last paragraph *	39,42	TECHNICAL FIELDS SEARCHED (Int. Cl. 4)
A		1-38, 40,41, 43-60	
A	-- NUCLEIC ACIDS RESEARCH, vol. 14, no. 12, June 1986, pages 4701-4718 IRL Press Ltd, Oxford, GB; J.M. PARDO et al.: "Cloning of the STA2 and SGA genes encoding glucoamylases in yeasts and regulation of their expression by the STA10 gene of Saccharomyces cerevisiae" * Abstract *	39,40	
X	-- EP - A - 0 185 327 (SUNTORY) * Page 10, line 29 - page 12, line 8; figure 6a *	39,45, 46,47	
Y	-- EP - A - 0 096 430 (GIST-BROCADES) * Page 14, lines 3-13 *	1-3, 7-9, 19,21, 22,28, 32,37, 47	
	--		

EPO Form 1505.J 08.78



DOCUMENTS CONSIDERED TO BE RELEVANT			CLASSIFICATION OF THE APPLICATION (Int. Cl.4)
Category	Citation of document with indication, where appropriate, of relevant passages	Relevant to claim	
Y	G. REIPEN et al.: "Non-selective transformation of <i>Saccharomyces cerevisiae</i> ", & CURR.GENET.1982, 6(3), 189-93 * Summary *	15	
Y	FOUNDATION FOR BIOTECHNICAL AND INDUSTRIAL FERMENTATION RESEARCH VI (1983), pages 209-228 A.M. SILLS et al.: "Genetic manipulation of amylolytic enzyme production by yeasts" * Page 210, paragraphs 2,3; page 220, lines 5-8; page 223 *	1-60	
A	EP - A 0 126 206 (CETUS) * Page 3, lines 1-9; page 5, line 32 - page 6, line 3, page 13, lines 18-23; line 31 - page 14, line 18; page 45, lines 15-17; page 46, line 1- page 47, line 6; claims 21-23 *	1-60	TECHNICAL FIELDS SEARCHED (Int. Cl.4)
P,X	WO - A - 86 07 091 (BIOTECHNICA) * Claims 1-5,9-16 *	1-4, 7-9, 19,21- 23,25, 27,28, 32,37, 39,40, 46-51, 53	
